Recommendations to Improve Children's Health Care Quality Measures

Background Report on the 2012 Process

Structured Abstract

Background: In 2009, the Children's Health Insurance Program Reauthorization Act (CHIPRA) (Public Law 111-3) directed the Secretary of the U.S. Department of Health and Human Services (HHS) to identify, by January 1, 2010, an initial core set of health care quality measures for voluntary use by Medicaid and the Children's Health Insurance Program (CHIP) programs. In addition, CHIPRA directed the Secretary to establish a CHIPRA Pediatric Quality Measures Program (PQMP). The purposes of the PQMP go beyond use by Medicaid and CHIP programs; the PQMP is to:

A. Improve and strengthen the initial core set of measures of health care quality established under CHIPRA.

B. Expand on existing pediatric quality measures used by public and private health care purchasers.

C. Increase the portfolio of evidence-based consensus pediatric quality measures available to public and private purchasers of children's health care services, providers, and consumers.

Measures developed under the PQMP are to be used to recommend changes to the initial core set beginning January 1, 2013, and annually thereafter. This document, measures recommended as changes to the initial core set (purpose A)

uses. CMS considered the SNAC recommendations and advised the Secretary of HHS on potential improvements to the initial core set.

Results: In all, 77 measures were submitted for consideration by the SNAC, 64 by the public and 13 by CHIPRA PQMP COEs.

Introduction

Initial Core Set

In 2009, the Children's Health Insurance Program Reauthorization Act (CHIPRA) (Public Law 111-3) amended Section 1139 of Title XX of the Social Security Act by adding Section 1139A

- x Updated periodically

- x Responsive to the child health needs, services, and domains of health care quality described in clauses (i), (ii), and (iii) of subsection (a)(6)(A)

CHIPRA also delineated a broad set of stakeholders to be included in the PQMP process (Section 1139A(b)(3)) and stipulated that grants and contracts could be used to develop, validate, and test a portfolio of pediatric quality measures (Section 1139A(b)(4)). Section 1139A(b)(5) required the Secretary of HHS to "publish recommended changes to the core measures described in subsection [(Section 1139A)] (a) that shall reflect the testing, validation, and consensus process for the development of pediatric quality measures described in subsection paragraphs (1) through (4)."

Measures developed under the PQMP are to be used to recommend changes to the initial core set beginning January 1, 2013, and annually thereafter. In this document, measures recommended as changes to the initial core set are referred to as additions to the core set or the improved core set. Measures meeting purposes (B) and (C) are referred to as measures for "other uses."

In response to the legislative directive to develop, validate, and test measures (Section 1139A(b)(4)) in March 2011, AHRQ and CMS awarded cooperative agreement grants to seven CHIPRA PQMP Centers of Excellence (COEs).[4] Each of these programs comprises multiple entities with diverse talents and expertise, including measurement capability, clinical knowledge, and the stakeholder groups of frontline providers, State Medicaid and CHIP programs, and patient representatives. Following a kickoff meeting with the COEs in April 2011, and consultation with CMS, the COEs were assigned an initial set of topics for me417.72 Tm ( )Tj 0. Tc 0 T

For SNAC scoring purposes, each measure's online CPCF form was expanded to include a function for scoring nine of the desirable measure attributes and three opportunities for "global scoring" (Table 1). In consideration of the limited time SNAC members had for scoring and the large numbers of measures on multiple attributes, scoring opportunities 1 through 9 were optional. SNAC members could score each of these separately (using the modified Delphi scoring approach on a scale of 1-9 described below) and calculate a combined global score, or they could simply read through the responses related to desirable measure attributes and more qualitatively assign a global score. In addition, SNAC members were asked to score on a scale of 1-5 whether a

Modified Delphi Process. A modified Delphi process using three rounds of scoring was used to determine the measures that would be considered for the improved core set and for other purposes. Delphi rounds I and II were completed by SNAC members prior to an in-person meeting held on September 12, 2012. The final (Delphi III) round to reach consensus on measures was conducted at the in-person meeting.

Delphi Round I. In Delphi Round I, the 50 publicly nominated measures that were extracted into the CPCF form were evaluated by the SNAC across a 4-week period. After reviewing a summary of the submitted scores during a webinar, the SNAC subsequently agreed on the following cutoff points for further consideration of a measure:

- If on Question 10, a measure received a median score of 7 or greater and had an interquartile range (IQR) of 2.25 or lower, indicating a low dispersion of scores, AND on Question 11 or Question 12 received a median score of 4 or greater and had an IQR of 1.25 or lower, the measure would be considered for the improved core set and/or other uses at the in-person meeting. In summary, measures that met this cutoff were considered to have received high scores.

- If the measure had a median score of greater than 7 for Question 10, regardless of the median score for Question 11 or 12, OR if the measure received a median score of 6 or greater and/or an IQR greater than 3 for Question 10, AND a median score of 4 or higher for Question 11 or 12, then the measure was to be voted on again in Delphi II.

- If a measure scored below the predetermined cutoff points in Delphi I and all preceding rounds, the measure would not receive further consideration in Delphi II or the in-person meeting.

taking all desirable measure attributes and information into consideration, as well as the results of votes for additions to the core set for the two other CHIPRA PQMP purposes.

In Delphi Round I, six measures scored highly enough to be voted on at the in-person meeting, 20 met criteria to be considered again in Round II, and 24 did not move forward. In Round II, of the 20 measures moved forward from Round I, five met criteria to be voted on for the improved core set and other purposes at the in-person meeting, 10 met criteria for Delphi III, and 18 did not move forward. Of the 13 measures submitted by the three CCOEs for initial consideration during Delphi Round I, two met criteria for voting for the improved core set at the in-person meeting, five met criteria for Delphi Round III, and six did not move forward. Of the 10 measures considered in Delphi III, two moved forward for consideration for the improved core set and other purposes, for a total of 13 measures for such consideration. The SNAC recommended five measures as additions to the core set. Finally, of the eight high-scoring measures that were not recommended as improvements to the core set, the SNAC recommended two measures for use by other private and public programs. A summary of measures recommended for the improved core set and for other purposes is presented in Table 2; each of the recommended measures is described in more detail in the following section

Table 2. SNAC-Recommended Children's Health Care Quality Measures for Additions to the Core Set in 2013 and for Two Other CHIPRA Purposes

Topic	Measure Name	Measure Submitter [1]
Recommended for voluntary use by Medicaid/CHIP and other purposes		
Duration of enrollment	Coverage in Medicaid and CHIP	CHOP
Duration of enrollment	Duration of newborn's first enrollment	CHOP
Child and adolescent immunizations	Human papillomavirus (HPV) vaccine for female adolescents	NCQA
Imaging	Recording radiation exposure from diagnostic computed tomography exams	St. Louis Children's Hospital
Asthma	Medication management for people with asthma	NCQA
Recommended for other purposes only		
Prenatal	Behavioral risk assessment	PMCoE-AMA PCPI
Substance use/abuse	Tobacco use and help with quitting among adolescents	NCINQ

Key: CHIP = Children's Health Insurance Program; CHIPRA = Children's Health Insurance Program Reauthorization Act; CHOP = Children's Hospital of Philadelphia (PQMPCOE); NCINQ = National Collaborative for Innovation in Quality Measurement (Pediatric Quality Measures Program [PQMP] Center of Excellence [COE]); NCQA = National Committee for Quality Assurance; PMCoE = Pediatric Measurement Center of Excellence (a PQMPCOE), of which the American Medical Association

COE. At the same time, CMS had to respond to the requirement in CHIPRA Section 402 for a measure to assess enrollment and retention data (including data with respect to continuity of coverage or duration of benefits).

This measure is flexible; the cohort can be defined at any time period of interest and can be tracked as long as data are available. Moreover, the definition of new enrollee can be varied. This measure is also sensitive to gaps in coverage and somewhat sensitive to trends in eligibility rates due to changes in state or nationwide economic downturns and upturns, which are periods of time that children tend to enroll and disenroll, respectively.

The SNAC supported recommending the CHOP COE measure primarily because of its feasibility—i.e., States can calculate coverage and enrollment from existing Medicaid and CHIP administrative data. Furthermore, it measures continuity of coverage by Medicaid or CHIP, rather than measuring coverage in each public program separately, thereby presenting a more complete picture of public insurance coverage. States may also find this measure valuable because the measure of coverage in Medicaid and CHIP could highlight State practices that reduce or increase administrative barriers to obtaining coverage in public programs, such as frequency of re-enrollment and documentation necessary for enrollment.

vaccination is most effective before a female is exposed to HPV, i.e., before they become sexually active. Thus, HPV vaccination is recommended at 11 and 12 years of age, as the median age of first sexual experience for Americans is 17 years, and studies suggest that 13 percent of girls initiate sexual activity before the age of 15.

This measure is part of the National Committee for Quality Assurance (NCQA) Health Effectiveness Data and Information Set (HEDIS) Health Plan measure set, which is a widely used instrument for health plan quality measurement and improvement. This measure complements the NCQA HEDIS Immunizations for Adolescents measure already in the initial core set, which assesses immunization status for meningococcal vaccine (MCV4), and one tetanus, diphtheria, and acellular pertussis vaccine (Tdap) or one tetanus, diphtheria toxoids vaccine (Td), by a child's 13th birthday, but does not include HPV vaccination. The HPV vaccine was added to the Advisory Committee on Immunization Practices (ACIP) vaccine schedule for adolescents in 2007.

The SNAC discussed how this measure reflects the current research on the importance of vaccination for HPV vaccine for the adult health outcome of preventing cervical cancer. SNAC members noted that by recommending this measure, the improved core set would reflect current, evidence-based practice guidelines.

Recording Radiation Exposure from Diagnostic Computerized Tomography (CT) Exams. This measure assesses the number of diagnostic computed tomography (CT) exams for which metrics of radiation exposure are documented in the electronic health record (EHR) as a proportion of CT scans performed at a facility. The measure requires including at least one metric of radiation exposure in the radiology report that describes a CT study's findings and interpretation. Suitable metrics include any of the following: Computed Tomography Dose Index Volume (CTDIvol), Dose Length Product (DLP), Size-Specific Dose Estimate (SSDE), or any other metric recommended by the American Association of Physicists in Medicine (AAPM).

The increasing numbers of CT scans performed in the pediatric population have raised concerns about the long-term risks of medical imaging to children.[18] SNAC members noted that this measure could encourage providers to measure and record the level of radiation exposure, providing a basis for documenting cumulative exposure, and argued that including this measure in the core set would encourage greater transparency about the radiation exposure children receive. However, they acknowledged this as an aspirational measure with potential challenges to implementation, especially since modality and reliability testing needs to be done. In addition, SNAC members and other participants in the SNAC meeting were aware that a COE had been assigned the task of developing a measure related to imaging.

Medication Management for People with Asthma. This measure assesses whether children 5 to 18 years of age with persistent asthma remain on their medications. A first rate examines the percentage of children who remained on an asthma controller medication prescription for at least 50 percent of their treatment period. The second rate examines those who remained on the medication for at least 75 percent of their treatment period. These rates are reported as the proportion of days covered by at least one prescription for an asthma controller medication, and the medication dispensed, during the measurement year.

This measure can be paired with the initial core set measure that calculates the annual percentage of asthma patients with one or more asthma-related emergency room visits. Over 7 million children under the age of 18 have asthma, placing it among the leading causes of hospitalization for children.[19] Appropriate medication management potentially could prevent a significant proportion of asthma-related costs (hospitalizations, emergency room visits, and missed work and school days).[20] In 2009, child hospitalizations for asthma as a principal diagnosis cost the nation $1.5 billion.[21]

The SNAC rationale for recommending this measure is that asthma is a common chronic pediatric health condition, and this measure focuses on disease management. Moreover, the SNAC considered that it is feasible to implement this measure in Medicaid and CHIP because it can be derived from existing administrative data.

Behavioral Risk Assessment During Prenatal Care. This measure assesses the percentage of patients, regardless of age, who gave birth during a 12-month period, were seen at least once for prenatal care, and received a behavioral health screening risk assessment. Risk assessments include the following screenings at the first prenatal visit: screening for depression, alcohol use, tobacco use, drug use, and intimate partner violence. This measure provides a mechanism to help identify pregnant women with drug, alcohol, or smoking problems, as well as depression and abuse, which may help prevent adverse neonatal outcomes. The adverse effects of alcohol, tobacco, and drugs on the fetus, including fetal alcohol syndrome and intrauterine growth restriction, are widely known.[22] In addition, women abused during pregnancy are more likely to be depressed, suicidal, and experience pregnancy complications and poor outcomes, including maternal and fetal death.[23]

SNAC members commented that the measure accounts for the mental and social factors involved in perinatal care. However, SNAC members were concerned that the measure relies on EHRs as its data source, and the use of EHRs is not yet universal among physicians. The measure would not be immediately feasible for use by all Medicaid and CHIP programs, according to the SNAC, which recommended the measure for other uses.

Tobacco Use and Help with Quitting Among Adolescents. This measure identifies the percentage of adolescents 12 to 20 years of age during the measurement year for whom tobacco use status was documented who received help with quitting if identified as a tobacco user. "Receiving help" is determined as documentation of any of the following: patient given advice to quit smoking or tobacco use, counseling on the benefits of quitting smoking or tobacco use, assistance with or referral to external smoking or tobacco cessation support programs, or current enrollment in smoking or tobacco use cessation program. The measure focuses on a clinical process (documentation of tobacco use and appropriate followup) that, if followed, has the potential to achieve a desirable clinical outcome (cessation of tobacco use, which can reduce the risk for a wide range of conditions known to be associated with tobacco use, such as asthma and lung cancer).

Smoking is a relevant issue in pediatric health care because over 2.6 million adolescents 18 years of age and younger are current tobacco users, with nearly 1 in 5 of all adolescents becoming current smokers before finishing high school.[24,25] Furthermore, of adults who smoke on a daily

basis, 82 percent reported trying their first cigarette before the age of 18, and 53 percent reported becoming daily smokers before the age of 18. The financial burden incurred from tobacco use is significant. From 2000 to 2004, annual expenditures (public and private) related to smoking were estimated to be $96 billion, and another $97 billion can be attributed to lost productivity each year.[26]

This measure was developed by the National Collaborative for Innovation in Quality Measurement (NCINQ) COE as part of its assignment to develop measures of the content of adolescent well-care. The COE submission provided, and the SNAC reviewed, results from measure testing that indicated that tobacco use status can be found in paper and electronic records, and that where found, documentation of followup ca(he)4( )-1- ou3,6 0 Tw 0 Tw 0

providers) and other sets of priorities (e.g., the National Quality Strategy,[29] the Institute of medicine [IOM] domains[30]). The SNAC also suggested refinements to the SNAC review process for subsequent years. For instance, to make the review process less burdensome, the SNAC suggested greater triaging of measures before assi10(as)N28.776neasures

References

[1] U.S. Congress and U.S. President. Public Law 111-3. Children's Health Insurance Program Reauthorization Act 2009; http://www.gpo.gov/fdsys/pkg/PLAW-111publ3/html/PLAW-111publ3.htm. Accessed Dec. 7, 2012.

[2] Agency for Healthcare Research and Quality. National Advisory Council on Healthcare Research and Quality. http://www.ahrq.gov/about/council.htm. Accessed Dec. 5, 2012.

[3] Centers for Medicare & Medicaid Services. Initial Core Set of Children's Health Care Quality Measures:

Appendix A: Members of the 2012 AHRQ Subcommittee to the National Advisory Council on Children's Health Quality Measures (SNAC)

Elizabeth Anderson*

Co-c/MCID /MCID 1 >>BDo on C2o, Pe,e2b, PondvP(e)-1oeh7s se El-13.56 Tw 12(ubc)-4(-8(m A)-ere

Glenn Flores, MD, FAAP *
Professor, Pediatrics and Public Health
University of Texas Southwestern
Director, Division of General Pediatrics
UT Southwestern Children's Medical Center of Dallas
Dallas, TX

Charles Gallia, PhD*
Senior Health Policy Advisor
Division of Medical Assistance Programs
Oregon Health Authority
Salem, OR

Paul Glassman, DDS, MA, MBA
Director
Pacific Center for Special Care
Professor, Department of Dental Care

Medicaid Program
Washington State Health Care Authority
Olympia, WA

Kirsten Thomsen, PA*
Commissioner
National Commission on Vision and Health
Alexandria, VA

Feliciano Yu, MD *
Associate Professor

Appendix B: Measure Scores by CHIPRA Category, Measure Name, and Scoring Round

CHIPRA Category/ Measure Topic/ Subtopic	Measure Number	Measure Name and Measure Submitter	Delphi Round I Global Summary Score (Q 10) (Median Score, Interquartile Range (IQR))	Delphi Round II Global Summary Score

			6,2	

Identification of Children

Child and Adolescent Immunizations	0	**Human papillomavirus vaccine for female adolescents (NCQA)**	8,2	Bypass	Bypass	Improved Core Set	Improved Core Set
Screening +/- Followup							
x	0	W h	5,3				
	0	-	6,2	5.5, 1.75			
	0	3 y	5,3				
	0	0 7					

[1] "Pass" indicates at least 70 percent of SNAC members scored measure as "3"= high, in Delphi III, and the measure was considered for recommendation for the core set and other purposes.

[2] If no more scores are shown, the measure was not included in subsequent Delphi rounds and/or not recommended for the c ore setfrfor r1 Tc 0.003