

References for Studies Included in Meta-analyses

Table 6.4. Meta-analysis (1996): Impact of having a tobacco use status identification system in place on rates of clinician intervention with their patients who smoke (n = 9 studies)

*Cohen SJ, Christen AG, Katz BP, et al. Counseling medical and dental patients about cigarette smoking: the impact of nicotine gum and chart reminders. *Am J Public Health* 1987;77(3):313-6.

Dietrich AJ, O'Connor GT, Keller A, et al. Cancer: improving early detection and prevention. A community practice randomised trial. *Br Med J* 1992;304(6835):687-91.

Fiore MC, Jorenby DE, Schensky AE, et al. Smoking status as the new vital sign: effect on assessment and intervention with patients who smoke. *Mayo Clin Proc* 1995;70(3):209-13.

Hahn DL, Berger MG. Implementation of a systematic health maintenance protocol in a private practice. *J Fam Pract* 1990;31(5):492-504.

Kottke TE, Solberg LI, Brekke ML, et al. A controlled trial to integrate smoking cessation advice into primary care practice: doctors helping smokers, round III. *J Fam Pract* 1992;34(6):701-8.

McIlvain HE, Susman JL, Manners MA, et al. Improving smoking cessation counseling by family practice residents. *J Fam Pract* 1992;34(6):745-9.

Robinson MD, Laurent SL, Little Jr JM. Including smoking status as a new vital sign: it works! *J Fam Pract* 1995;40(6):556-61.

Strecher VJ, O'Malley MS, Villagra VG, et al. Can residents be trained to counsel patients about quitting smoking? Results from a randomized trial. *J Gen Intern Med* 1991;6(1):9-17.

* Article contributed two studies to the meta-analysis

Table 6.5. Meta-analysis (1996): Impact of having a tobacco use status identification system in place on abstinence rates among patients who smoke (n = 3 studies)

*Cohen SJ, Christen AG, Katz BP, et al. Counseling medical and dental patients about cigarette smoking: the impact of nicotine gum and chart reminders. *Am J Public Health* 1987;77(3):313-6.

Strecher VJ, O'Malley MS, Villagra VG, et al. Can residents be trained to counsel patients about quitting smoking? Results from a randomized trial. *J Gen Intern Med* 1991;6(1):9-17

* Article contributed two studies to the meta-analysis

Table 6.7. Meta-analysis (1996): Effectiveness of and estimated abstinence rates for advice to quit by a physician (n = 7 studies)

Demers RY, Neale AV, Adams R, et al. The impact of physicians' brief smoking cessation counseling: a MIRNET study. *J Fam Pract* 1990;31(6):625-9.

Jamrozik K, Vessey M, Fowler G, et al. Controlled trial of three different antismoking interventions in general practice. *Br Med J* 1984;288(6429):1499-1503.

Page AR, Walters DJ, Schlegel RP, et al. Smoking cessation in family practice: the effects of advice and nicotine chewing gum prescription. *Addict Behav* 1986;11(4):443-6.

Russell MA, Wilson C, Taylor C, et al. Effect of general practitioners' advice against smoking. *Br Med J* 1979;2(6184):231-5.

Russell MAH, Merriman R, Stapleton J, et al. Effect of nicotine chewing gum as an adjunct to general practitioners' advice against smoking. *Br Med J* 1983;287(6407):1782-5.

Slama K, Redman S, Perkins J, et al. The effectiveness of two smoking cessation programmes for use in general practice: a randomised clinical trial. *Br Med J* 1990;300(6741):1707-9.

Stewart PJ, Rosser WW. The impact of routine advice on smoking cessation from family physicians. *Can Med Assoc J* 1982;126(9):1051-4.

**Table 6.8. Meta-analysis (2000):
Effectiveness of and estimated abstinence rates for various intensity levels of session length (n = 43 studies)**

Barbarin OA. Comparison of symbolic and overt aversion in the self-control of smoking. *J Consult Clin Psychol* 1978;46(6):1569-71.

British Thoracic Society. Smoking cessation in patients: two further studies by the British Thoracic Society. *Thorax* 1990;45(11):835-40.

BronsoBC89 0 T/P <</MCID 5 >>BDC -0.002 TcoB.002 Tc 0.s/7p(i)3(-g7t)-1(s)-8(:)]TJ 0 Tc 0 Tw 3.783 0 Td ()(ur)-6(t)

Killen JD, Fortmann SP, Newman B, et al.
Evaluation of a treatment approach combining
nicotine gum with self-guided behavioral
treatments for smoking relapse prevention. *J
Consult Clin Psychol* 1990;58(1):85-92.

Lambe R, Osier C, Franks P. A randomized
controlled trial of hypnotherapy for smoking
cessation. *J Fam Pract* 1986;22(1):61-5.

Lewis SF, Piasecki TM, Fiore MC, et al.
Transdermal nicotine replacement for
hospitalized patients: a randomized clinical trial.
Prev Med 1998;27(2):296-303.

Ockene JK, Kristeller J, Goldberg R, et al.
Increasing the efficacy of physician-delivered
smoking interventions: a randomized clinical
trial. *J Gen Intern Med* 1991;6(1):1-8.

Ockene JK, Kristeller JL, Goldberg R, et al.
Smoking cessation and severity of disease: the
coronary artery smoking intervention study.
Health Psychol 1992;11(2):119-26.

Ossip-a

Weissfeld JL, Holloway JL. Treatment for cigarette smoking in a Department of Veterans Affairs outpatient clinic. *Arch Intern Med* 1991;151(5):973-7.

Williams JM, Hall DW. Use of single session hypnosis for smoking cessation. *Addict Behav* 1988;13(2):205-8.

Wilson D, Wood G, Johnston N, et al. Randomized clinical trial of supportive follow-up for cigarette smokers in a family practice. *Can Med Assoc J* 1982;126(2):127-9.

Windsor RA, Lowe JB, Bartlett EE. The effectiveness of a worksite self-help smoking cessation program: a randomized trial. *J Behav Med* 1988;11(4):407-21.

Zhu SH, Stretch V, Balabanis M, et al. Telephone counseling for smoking cessation: effects of single-session and multiple-session interventions. *J Consult Clin Psychol* 1996;64(1):202-11.

Table 6.9. Meta-analysis (2000): Effectiveness of and estimated abstinence rates for total amount of contact time (n = 35 studies)

Barbarin OA. Comparison of symbolic and overt aversion in the self-control of smoking. *J Consult Clin Psychol* 1978;46:1569-71.

British Thoracic Society. Smoking cessation in patients: two further studies by the British Thoracic Society. *Thorax* 1990;45(11):835-40.

Bronson DL, Flynn BS, Solomon LJ, et al. Smoking cessation counseling during periodic health examinations. *Arch Intern Med* 1989;149:1653-6.

Burling TA, Marotta J, Gonzalez R, et al. Computerized smoking cessation program for the worksite: treatment outcome and feasibility. *J Consult Clin Psychol* 1989;57:619-22.

Cottraux JA, Harf R, Boissel JP, et al. Smoking cessation with behaviour therapy or acupuncture—A controlled study. *Behav Res Ther* 1983;21:417-24.

Danaher BG, Jeffery RW, Zimmerman R, et al. Aversive smoking using printed instructions and audiotape adjuncts. *Addict Behav* 1980;5:353-8.

Fagerberg B, Wikstrand J, Berglund G, et al. Mortality rates in treated hypertensive men with additional risk factors are high but can be reduced: a randomized intervention study. *Am J Hypertens* 1998;11:14-22.

Fortmann SP, Killen JD, Telch MJ, et al. Minimal contact treatment for smoking cessation. A placebo controlled trial of nicotine polacrilex and self-directed relapse prevention: initial results of the Stanford stop smoking project. *JAMA* 1988;260:1575-80.

Gillams J, Lewith GT, Machin D. Acupuncture and group therapy in stopping smoking. *Practitioner* 1984;228:341-4.

Hall SM, Tunstall CD, Ginsberg D, et al. Nicotine gum and behavioral treatment: a placebo controlled trial. *J Consult Clin Psychol* 1987;55:603-5.

Hilleman DE, Mohiuddin SM, Delcore MG, et al. Randomized, controlled trial of transdermal clonidine for smoking cessation. *Ann Pharmacother* 1993;27:1025-8.

Hollis JF, Lichtenstein E, Vogt TM, et al. Nurse-assisted counseling for smokers in primary care. *Ann Intern Med* 1993;118:521-5.

Hyman GJ, Stanley RO, Burrows GD, et al. Treatment effectiveness of hypnosis and behaviour therapy in smoking cessation: a methodological refinement. *Addict Behav* 1986;11:355-65.

Jorenby DE, Smith SS, Fiore MC, et al. Varying nicotine patch dose and type of smoking cessation counseling. *JAMA* 1995;274:1347-52.

Killen JD, Fortmann SP, Newman B, et al. *Evalu*:1347-

Lando HA. Effects on preparation, experimenter contact, and a maintained reduction alternative on a broad-spectrum program for eliminating smoking. *Addict Behav* 1981;6:123-33.

Lando HA. Successful treatment of smokers with a broad-spectrum behavioral approach. *J Consult Clin Psychol* 1977;45:361-6.

Lando HA, K reduci12(o)]1(t)ch

Lando HA. Effects on preparation, experimenter contact, and a maintained reduction alternative on a broad-spectrum program for eliminating smoking. *Addict Behav* 1981;6(2):123-33.

Lando HA. Successful treatment of smokers with a broad-spectrum behavioral approach. *J Consult Clin Psychol* 1977;45(3):361-6.

Lando HA, Rolnick S, Klevan D, et al. Telephone support as an adjunct to transdermal nicotine in smoking cessation. *Am J Public Health* 1997;87(10):1670-74.

Lewis SF, Piasecki TM, Fiore MC, et al. Transdermal nicotine replacement for hospitalized patients: a randomized clinical trial. *Prev Med* 1998;27(2):296-303.

Lifrak P, Gariti P, Alterman AI, et al. Results of two levels of adjunctive treatment used with the nicotine patch. *Am J Addict* 1997;6(2):93-8.

Ockene JK, Kristeller J, Goldberg R, et al. Increasing the efficacy of physician-delivered smoking interventions: a randomized clinical trial. *J Gen Intern Med* 1991;6(1):1-8.

Ossip-Klein DJ, Carosella AM, Krusch DA. Self-help interventions for older smokers. *Tob Control* 1997;6(3):188-93.

Patel C, Marmot MG, Terry DJ, et al. Trial of relaxation in reducing coronary risk: four year follow up. *Br Med J* 1985;290(6475):1103-6.

Pederson LL, Scrimgeour WG, Lefcoe NM. Comparison of hypnosis plus counseling, counseling alone, and hypnosis alone in a community service smoking withdrawal program. *J Consult Clin Psychol* 1975;43(6):920.

Pederson LL, Wanklin JM, Lefcoe NM. The effects of counseling on smoking cessation among patients hospitalized with chronic obstructive pulmonary disease: a randomized clinical trial. *Int J Addict* 1991;26(1):107-19.

Poole AD, Sanson-Fisher RW, German GA. The rapid-smoking technique: therapeutic effectiveness. *Behav Res Ther* 1981;19(5):389-97.

Rabkin SW, Boyko E, Shane F, et al. A randomized trial comparing smoking cessation programs utilizing behaviour modification, health education or hypnosis. *Addict Behav* 1984;9:157-73.

Rand CS, Stitzer ML, Bigelow GE, et al. The effects of contingent payment and frequent workplace monitoring on smoking abstinence. *Addict Behav* 1989;14(2):121-8.

Reid DD, Brett GZ, Hamilton PJ, et al. Cardiorespiratory disease and diabetes among middle-aged male civil servants. A study of screening and intervention. *Lancet* 1974;1(7856):469-73.

Rigotti NA, Arnsten JH, McKool KM, et al. Efficacy of a smoking cessation program for hospital patients. *Arch Intern Med* 1997;157(22):2653-60.

Segnan N, Ponti A, Battista RN, et al. A randomized trial of smoking cessation interventions in general practice in Italy. *Cancer Causes Control* 1991;2(4):239-46.

*Sivarajan ES, Newton KM, Almes MJ, et al. Limited effects of outpatient teaching and c

**Table 6.11. Meta-analysis (2000):
Effectiveness of and estimated abstinence
rates for interventions delivered by different
types of clinicians (n = 29 studies)**

Brandon TH, Zelman DC, Baker TB. Effects of maintenance sessions on smoking relapse: delaying the inevitable? *J Consult Clin Psychol* 1987;55(5):780-2.

British Thoracic Society. Comparison of four methods of smoking withdrawal in patients with smoking related diseases. *Br Med J* 1983;286(6365):595-7.

Carlsson R, Lindberg G, Westin L, et al. Influence of coronary nursing management follow up on lifestyle after acute myocardial infarction. *Heart* 1997;77(3):256-9.

Cottraux JA, Harf R, Boissel JP, et al. Smoking cessation with behaviour therapy or acupuncture: a controlled study. *Behav Res Ther* 1983;21(4):417-24.

Curry SJ, McBride C, Grothaus LC, et al. A randomized trial of self-help materials, personalized feedback, and telephone counseling with nonvolunteer smokers. *J Consult Clin Psychol* 1995;63(6):1005-14.

Danaher BG, Jeffery RW, Zimmerman R, et al. Aversive smoking using printed instructions and audiotape adjuncts. *Addict Behav* 1980;5:353-8.

Davies BL, Matte-Lewis L, O'Connor AM, et al. Evaluation of the "time to quit" self-help smoking cessation program. *Can J Public Health* 1992;83(1):19-23.

Fortmann SP, Killen JD, Telch MJ, et al. Minimal contact treatment for smoking cessation. A placebo controlled trial of nicotine polacrilex and self-

**Table 6.12. Meta-analysis (2000):
Effectiveness of and estimated abstinence**

Rabkin SW, Boyko E, Shane F, et al. A randomized trial comparing smoking cessation programs utilizing behaviour modification, health education or hypnosis. *Addict Behav* 1984;9:157-73.

Reid DD, Brett GZ, Hamilton PJ, et al. Cardiorespiratory disease and diabetes among middle-aged male civil servants. A study of screening and intervention. *Lancet* 1974;1(7856):469-73.

Slama K, Redman S, Perkins J, et al. The effectiveness of two smoking cessation programmes for use in general practice: a randomised clinical trial. *Br Med J* 1990;300(6741):1707-9.

Stevens VJ, Severson H, Lichtenstein E, et al. Making the most of a teachable moment: a smokeless tobacco cessation intervention in the dental office. *Am J Public Health* 1995;85(2):231-5.

Stewart PJ, Rosser WW. The impact of routine advice on smoking cessation from family physicians. *Can Med Assoc J* 1982;126(9):1051-4.

Thompson B, Kinne S, Lewis FM, et al. Randomized telephone smoking-intervention trial initially directed at blue-collar workers. *J Natl Cancer Inst Monogr* 1993;(14):105-12.

Tonnesen P, Mikkelsen K, Markholst C, et al. Nurse-conducted smoking cessation with minimal intervention in a lung clinic: a randomized controlled study. *Eur Respir J* 1996;9(11):2351-5.

Weissfeld JL, Holloway JL. Treatment for cigarette smoking in a Department of Veterans Affairs outpatient clinic. *Arch Intern Med* 1991;151(5):973-7.

Williams JM, Hall DW. Use of single session hypnosis for smoking cessation. *Addict Behav* 1988;13(2):205-8.

Zhu SH, Stretch V, Balabanis M, et al. Telephone counseling for smoking cessation: effects of single-session and multiple-session interventions. *J Consult Clin Psychol* 1996;64(1):202-11.

Stewart PJ, Rosser WW. The impact of routine advice on smoking cessation from family physicians. *Can Med Assoc J* 1982;126(9):1051-4.

Taylor CB, Houston-Miller N, Killen JD, et al. Smoking cessation after acute myocardial infarction: effects of a nurse-managed intervention. *Ann Intern Med* 1990;113(2):118-23.

Tonnesen P, Mikkelsen K, Markholst C, et al. Nurse-conducted smoking cessation with minimal intervention in a lung clinic: a randomized controlled study. *Eur Respir J* 1996;9(11):2351-5.

Vetter NJ, Ford D. Smoking prevention among people aged 60 and over: a randomized controlled trial. *Age Ageing* 1990;19(3):164-6.

Weissfeld JL, Holloway JL. Treatment for cigarette smoking in a Department of Veterans Affairs outpatient clinic. *Arch Intern Med* 1991;151(5):973-7.

Williams JM, Hall DW. Use of single session hypnosis for smoking cessation. *Addict Behav* 1988;13(2):205-8.

Zhu SH, Stretch V, Balabanis M, et al. Telephone counseling for smoking cessation: effects of single-session and multiple-session interventions. *J Consult Clin Psychol* 1996;64(1):202-11.

**Table 6.13. Meta-analysis (2000):
Effectiveness of and estimated abstinence rates for various types of format (n = 58 studies)**

Barbarin OA. Comparison of symbolic and overt aversion in the self-control of smoking. *J Consult Clin Psychol* 1978;46(6):1569-71.

Brandon TH, Zelman DC, Baker TB. Effects of maintenance sessions on smoking relapse: delaying the inevitable? *J Consult Clin Psychol* 1987;55(5):780-

etu tmokie av(n(i)our6(:)--8(e(h8(u)-e(i)-1ap9(a)-12(yA83()-or6(:)

Gillams J, Lewith GT, Machin D. Acupuncture and group therapy in stopping smoking. *Practitioner* 1984;228(1389):341-4.

Glasgow RE, Schafer L, O'Neill HK. Self-help books and amount of therapist contact in smoking cessation programs. *J Consult Clin Psychol* 1981;49(5):659-67.

Gritz ER, Berman BA, Bastani R, et al. A randomized trial of a self-help smoking cessation intervention in a nonvolunteer female population: testing the limits of the public health model. *Health Psychol* 1992;11(5):280-9.

Hall SM, Munoz RF, Reus VI, et al. Mood management and nicotine gum in smoking treatment: a therapeutic contact and placebo-controlled study. *J Consult Clin Psychol* 1996;64(5):1003-9.

Hall SM, Tunstall CD, Vila KL, et al. Weight gain prevention and smoking cessation: cautionary findings. *Am J Public Health* 1992;82(6):799-803.

Hilleman DE, Mohiuddin SM, Delcore MG, et al. Randomized, controlled trial of transdermal clonidine for smoking cessation. *Ann Pharmacother* 1993;27(9):1025-8.

Hollis JF, Lichtenstein E, Vogt TM, et al. Nurse-assisted counseling for smokers in primary care. *Ann Intern Med* 1993;118(7):521-5.

Hughes GH, Hymowitz N, Ockene JK, et al. The multiple risk f

d trial3(l163(C)-3()-1(r)-6(ans)-8(d3(l163(m)-24(ac)-8(6 0 -)Tj --)Tj -e2(993)-12(Tw 2(

Ockene J, Kristeller JL, Goldberg R, et al. Smoking cessation and severity of disease: the coronary artery smoking intervention study. *Health Psychol* 1992;11(2):119-26.

Ockene JK, Kristeller J, Goldberg R, et al. Increasing the efficacy of physician-delivered smoking interventions: a randomized clinical trial. *J Gen Intern Med* 1991;6(1):1-8.

OSSIP-Klein DJ, Carosella AM, Krusch DA. Self-help interventions for older smokers. *Tob Control* 1997;6(3):188-93.

Patel C, Marmot MG, Terry DJ, et al. Trial of relaxation in reducing coronary risk: four year follow up. *Br Med J* 1985;290(6475):1103-6.

Pederson LL, Scrimgeour WG, Lefcoe NM. Comparison of hypnosis plus counseling, counseling alone, and hypnosis alone in a community service smoking withdrawal program. *J Consult Clin Psychol* 1975;43(6):920.

Pederson LL, Wanklin JM, Lefcoe NM. The effects of counseling on smoking cessation among patients hospitalized with chronic obstructive pulmonary disease: a randomized clinical trial. *Int J Addict* 1991;26(1):107-

Hall SM, Munoz RF, Reus VI, et al. Mood management and nicotine gum in smoking treatment: a therapeutic contact and placebo-controlled study. J Consult Clin Psychol 1996;64(5):1003-9.

Hall SM, Tunstall CD, Vila KL, et al. Weight gain prevention and smoking cessation: cautionary findings. Am J Public Health 1992;82(6):799-803.

Hilleman DE, Mohiuddin SM, Delco2(s)-81 >)-7 II16.229 ()Tj

II16.229 ()Tj

he Pu(s)-329 (,p(s)-3e(s)-)Tj -0.002 Tc 0.2.86 0 Td (r2 Tw -4.052 -2.5665Tj E0.34-1(i)-9(ois TTc es)-)-

Hall SM, Reus VI, Munoz RF, et al. Nortriptyline and cognitive-behavioral therapy in the treatment of cigarette smoking. *Arch Gen Psychiatry* 1998;55(8):683-90.

Hall SM, Tunstall CD, Ginsberg D, et al. Nicotine gum and behavioral treatment: a placebo-controlled trial. *J Consult Clin Psychol* 1987;55(4):603-5.

Hall SM, Tunstall CD, Vila KL, et al. Weight gain prevention and smoking cessation: cautionary findings. *Am J Public Health* 1992;82(6):799-803.

Hollis JF, Lichtenstein E, Vogt TM, et al. Nurse-assisted counseling for smokers in primary care. *Ann Intern Med* 1993;118(7):521-5.

Hughes GH, Hymowitz N, Ockene JK, et al. The multiple risk factor intervention trial (MRFIT)ea -0.002 Tc 0.002 T Ann n Meion

Slovinec D'Angelo ME, Reid RD, Hotz S, et al. Is stress management training a useful addition to

Aubin HJ, Lebargy F, Berlin I, et al. Efficacy of bupropion and predictors of successful outcome in a sample of French smokers: a randomized placebo-controlled trial. *Addiction* 2004;99:1206-18.

Blondal T, Gudmundsson LJ, Olafsdottir I, et al. Nicotine nasal spray with nicotine patch for smoking cessation: randomised trial with six year follow up. *Br Med J* 1999;318:285-8.

Bohadana A, Nilsson F, Rasmussen T, et al. Nicotine inhaler and nicotine patch as a combination therapy for smoking cessation: a randomized, double-blind, placebo-controlled trial. *Arch Intern Med* 2000;160:3128-34.

Cinciripini PM, Tsoh JY, Wetter DW, et al. Combined effects of venlafaxine, nicotine replacement, and brief counseling on smoking cessation. *Exp Clin Psychopharmacol* 2005;13(4):282-92.

Collins BN, Wileyto EP, Patterson F, et al. Gender differences in smoking cessation in a placebo-controlled trial of bupropion with behavioral counseling. *Nicotine Tob Res* 2004;6(1):27-37.

Cooper TV, Klesges RC, Debon MW, et al. A placebo controlled randomized trial of the effects of phenylpropanolamine and nicotine gum on cessation rates and postcessation weight gain in women. *Addict Behav* 2005;30(1):61-75.

Covey LS, Glassman AH, Stetner F, et al. A randomized trial of sertraline as a cessation aid for smokers with a history of major depression. *Am J Psychiatry* 2002;159:1731-7.

Croghan GA, Sloan JA, Croghan IT, et al. Comparison of nicotine patch alone versus nicotine nasal spray alone versus a combination for treating smokers: a minimal intervention,

Glassman AH, Covey LS, Dalack GW

Jorenby DE, Hays JT, Rigotti NA, et al. Efficacy of varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs placebo or sustained-release bupropion for smoking cessation: a randomized controlled trial. *JAMA* 2006;296(1):56-63.

Jorenby DE, Leischow SJ, Nides MA, et al. A controlled trial of sustained-release bupropion, a nicotine patch, or both for smoking cessation. *New Eng J Med* 1999;340(9):685-91.

Joseph AM, Norman SM, Ferry LH, et al. The safety of transdermal nicotine as an aid to smoking cessation in patients with cardiac disease. *N Eng J Med* 1996;335(24):1792-8.

Killen JD, Fortmann SP, Davis L, et al. Nicotine patch and self-help video for cigarette smoking cessation. *J Consult Clin Psychol* 1997;65:663-72.

Killen JD, Fortmann SP, Murphy Jr GM, et al. Extended treatment with bupropion SR for cigarette smoking cessation. *J Consult Clin Psychol* 2006;74:286-94.

Killen JD, Fortmann SP, Newman B, et al. Evaluation of a treatment approach combining nicotine gum with self-guided behavioral treatments for smoking relapse prevention. *J Consult Clin Psychol*

*Schneider NG, Jarvik ME, Forsythe AB, et al. Nicotine gum in smoking cessation: a placebo-controlled, double-blind trial. *Addict Behav* 1983;8:253-61.

Schneider NG, Olmstead R, Mody FV, et al. Efficacy of a nicotine nasal spray in smoking cessation: a placebo-controlled, double-blind trial. *Addiction* 1995;90:1671-82.

Schneider NG, Olmstead R, Nilsson F, et al. Efficacy of a nicotine inhaler in smoking cessation: a double-blind, placebo-controlled trial. *Addiction* 1996;91:1293-306.

Simon JA, Duncan C, Carmody TP, et al. Bupropion for smoking cessation: a randomized trial. *Arch Intern Med* 2004;164:1797-803.

*Sonderskov J, Olsen J, Sabroe S, et al. Nicotine patches in smoking cessation: a randomized trial among over-the-counter customers in Denmark. *Am J Epidemiol* 1997;145:309-18.

Stapleton JA, Russell MA, Feyerabend C, et al. Dose effects and predictors of outcome in a randomized trial of transdermal nicotine patches in general practice. *Addiction* 1995;90:31-42.

Sutherland G, Stapleton JA, Russell MA, et al. Randomised controlled trial of nasal nicotine spray in smoking cessation. *Lancet* 1992;340:324-9.

Swanson NA, Burroughs CC, Long MA, et al. Controlled trial for smoking cessation in a Navy shipboard population using nicotine patch, sustained-release bupropion, or both. *Mil Med* 2003;168:830-4.

Tashkin D, Kanner R, Bailey W, et al. Smoking cessation in patients with chronic obstructive pulmonary disease: a double-blind, placebo-controlled, randomised trial. *Lancet* 2001;357(9268):1571-5.

Tonnesen P, Mikkelsen KL. Smoking cessation with four nicotine replacement regimes in a lung clinic. *Eur Respir J* 2000;16:717-22.

Tonnesen P, Norregaard J, Mikkelsen K, et al. A double-blind trial of a nicotine inhaler for smoking cessation. *JAMA* 1993;269:1268-71.

Tonnesen P, Norregaard J, Simonsen K, et al. A double-blind trial of a 16-hour transdermal nicotine patch in smoking cessation. *N Engl J Med* 1991;325:311-5.

Tonnesen P, Paoletti P, Gustavsson G, et al. Higher dosage nicotine patches increase one-year smoking cessation rates: results from the European CEASE trial. *Eur Respir J* 1999;13(2):238-46.

Tonnesen P, Tonstad S, Hjalmarson A, et al. A multicentre, randomized, double-blind, placebo-controlled, 1-year study of bupropion SR for smoking cessation. *J Intern Med* 2003;254:184-92.

Tonstad S, Farsang ^{bl}
C

Davidson M, Epstein M, Burt R, et al. Efficacy and safety of an over-the-counter transdermal nicotine patch as an aid for smoking cessation. *Arch Fam Med* 1998;7:569-74.

Evins AE, Cather C, Deckersbach T, et al. A double-blind placebo-controlled trial of bupropion sustained-release for smoking cessation in schizophrenia. *J Clin Psychopharmacol* 2005;25:218-25.

Evins AE, Mays VK, Rigotti NA, et al. A pilot trial of bupropion added to cognitive behavioral therapy for smoking cessation in schizophrenia. *Nicotine Tob Res* 2001;3:397-403.

Fee WM, Stewart MJ. A controlled trial of nicotine chewing gum in a smoking withdrawal clinic. *Practitioner* 1982;226:148-151.

*Fiore MC, Kenford SL, Jorenby DE, et al. Two studies of the clinical effectiveness of the nicotine patch with different counseling treatments. *Chest* 1994;105:524-33.

Fortmann SP, Killen JD, Telch MJ, et al. Minimal contact treatment for smoking cessation. A placebo controlled trial of nicotine polacrilex and self-directed relapse prevention: initial results of the Stanford stop smoking project. *JAMA* 1988;260:1575-80.

George TP, Vessicchio JC, Termine A, et al. A placebo controlled trial of bupropion for smoking cessation in schizophrenia. *Biol Psychiatry* 2002;52:53-61.

Glassman AH, Covey LS, Dalack GW, et al. Smoking cessation, clonidine, and vulnerability to nicotine among dependent smokers. *Clin Pharmacol Ther* 1993;54:670-9.

Glassman AH, Stetner F, Walsh BT, et al. Heavy smokers, smoking cessation, and clonidine: results of a double-blind, randomized trial. *JAMA* 1988;259:2863-6.

Gonzales D, Rennard SI, Nides M, et al. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs sustained-release bupropion and placebo for smoking cessation: a randomized controlled trial. *JAMA* 2006;296:47-55.

Gonzales DH, Nides MA, Ferry LH, et al. Bupropion SR as an aid to smoking cessation in smokers treated previously with bupropion: a randomized placebo-controlled study. *Clin Pharmacol Ther* 2001;69:438-44.

Gourlay SG, Forbes A, Marriner T, et al. Double blind trial of repeated treatment with transdermal nicotine for relapsed smokers. *Br Med J* 1995;311:363-6.

Haggstram FM, Chatkin JM, Sussenbach-Vaz E, et al. A controlled trial of nortriptyline, sustained-release bupropion and placebo for smoking cessation: preliminary results. *Pulm Pharmacol Ther* 2006;19:205-9.

Hall SM, Humfleet GL, Reus VI, et al. Extended nortriptyline and psychological treatment for cigarette smoking. *Am J Psychiatry* 2004;161:2100-7.

Hall SM, Humfleet GL, Reus VI, et al. Psychological intervention and antidepressant treatment in smoking cessation. *Arch Gen Psychiatry* 2002;59:930-6.

Hall SM, Munoz RF, Reus VI, et al. Mood management and nicotine gum in smoking treatment: a therapeutic contact and placebo-controlled study. *J Consult Clin Psychol* 1996;64:1003-9.

Hall SM, Tunstall CD, Ginsberg D, et al. Nicotine gum and behavioral treatment: a placebo controlled trial. *J Consult Clin Psychol* 1987;55:603-5.

Hays JT, Croghan IT, Schroeder DR, et al. Over-the-counter nicotine patch therapy for smoking cessation: results from randomized, double-blind, placebo-controlled, and open label trials. *Am J Public Health* 1999;89:1701-7.

Herrera N, Franco R, Herrera L, et al. Nicotine gum, 2 and 4 mg, for nicotine dependence. A double-blind placebo-controlled trial within a behavior modification support program. *Chest* 1995;108:447-51.

Hjalmarson A, Franzon M, Westin A, et al. Effect of nicotine nasal spray on smoking cessation. A randomized, placebo-controlled, double-blind study. *Arch Intern Med* 1994;154:2567-72.

Hjalmarson A, Nilsson F, Sjostrom L, et al. The nicotine inhaler in smoking cessation. *Arch Intern Med* 1997;157:1721-8.

Holt S, Timu-Parata C, Ryder-Lewis S, et al. Efficacy of bupropion in the indigenous Maori population in New Zealand. *Thorax* 2005;60:120-3.

Hughes JR, Lesmes GR, Hatsukami DK, et al. Are higher doses of nicotine replacement more effective for smoking cessation? *Nicotine Tob Res* 1999;1:169-74.

Hughes JR, Novy P, Hatsukami DK, et al. Efficacy of nicotine patch in smokers with a history of alcoholism. *Alcohol Clin Exp Res* 2003;27:946-54.

Hurt RD, Dale LC, Fredrickson PA, et al. Nicotine patch therapy for smoking cessation combined with physician advice and nurse follow-up. One-year outcome and percentage of nicotine replacement. *JAMA* 1994;271:595-600.

Hurt RD, Sachs DP, Glover ED, et al. A comparison of sustained-release bupropion and placebo for smoking cessation. *N Eng J Med* 1997;337:1195-202.

Jensen EJ, Schmidt E, Pedersen B, et al. The effect of nicotine, silver acetate, and placebo chewing gum on the cessation of smoking. The influence of smoking type and nicotine dependence. *Int J Addict* 1991;26:1223-31.

Jorenby DE, Hays JT, Rigotti NA, et al. Efficacy of varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs placebo or sustained-release bupropion for smoking cessation: a randomized controlled trial. *JAMA* 2006;296:56-63.

Jorenby DE, Leischow SJ, Nides MA, et al. A controlled trial of sustained-release bupropion, a nicotine patch, or both for smoking cessation. *N Engl J Med* 1999;340:685-91.

Joseph AM, Norman SM, Ferry LH, et al. The safety of transdermal nicotine as an aid to smoking cessation in patients with cardiac disease. *N Eng J Med* 1996;335:1792-8.

Killen JD, Fortmann SP, Davis L, et al. Nicotine patch and self-help video for cigarette smoking cessation. *J Consult Clin Psychol* 1997;65:663-72.

Killen JD, Fortmann SP, Murphy Jr GM, et al. Extended treatment with bupropion SR for cigarette smoking cessation. *J Consult Clin Psychol* 2006;74:286-94.

Killen JD, Fortmann SP, Newman B, et al. Evaluation of a treatment approach combining nicotine gum with self-guided behavioral treatments for smoking relapse prevention. *J Consult Clin Psychol* 1990;58:85-92.

Killen JD, Fortmann SP, Schatzberg AF, et al. Nicotine patch and paroxetine for smoking cessation. *J Consult Clin Psychol* 2000;68:883-9.

Kornitzer M, Boutsen M, Dramaix M, et al. Combined use of nicotine patch and gum in smoking cessation: a placebo-controlled clinical trial. *Prev Med* 1995;24:41-7.

Leischow SJ, Nilsson F, Franzon M, et al. Efficacy of the nicotine inhaler as an adjunct to smoking cessation. *Am J Health Behav* 1996;20(5):364-71.

Lewis SF, Piasecki TM, Fiore MC, et al. Transdermal nicotine replacement for hospitalized patients: a randomized clinical trial. *Prev Med* 1998;27:296-303.

Myles PS, Leslie K, Angliss M, et al. Effectiveness of bupropion as an aid to stopping smoking before elective surgery: a randomised controlled trial. *Anaesthesia* 2004;59:1053-8.

Niaura R, Spring B, Borrelli B, et al. Multicenter trial of fluoxetine as an adjunct to behavioral smoking cessation treatment. *J Consult Clin Psychol* 2002;70:887-96.

Nides M, Oncken C, Gonzales D, et al. Smoking cessation with varenicline, a selective alpha4beta2 nicotinic receptor partial agonist: results from a 7-week, randomized, placebo- and bupropion-controlled trial with 1-year follow-up. *Arch Intern Med* 2006;166:1561-8.

Oncken C, Gonzales D, Nides M, et al. Efficacy and safety of the novel selective nicotinic acetylcholinergic antagonist, 3-(4-(2,6-dichlorophenyl)-5-methylpyridin-3-yl)pyridine (T4(α)-15(10H)-8(e)D

Tonnesen P, Norregaard J, Simonsen K, et al. A double-blind trial of a 16-hour transdermal nicotine patch in smoking cessation. *N Engl J Med* 1991;325:311-5.

Tonnesen P, Paoletti P, Gustavsson G, et al. Higher dosage nicotine patches increase one-year smoking cessation rates: results from the European CEASE trial. *Eur Respir J* 1999;13:238-46.

Tonnesen P, Tonstad S, Hjalmarson A, et al. A multicentre, randomized, double-blind, placebo-controlled, 1-year study of bupropion SR for smoking cessation. *J Intern Med* 2003;254:184-92.

Tonstad S, Farsang C, Klaene G, et al. Bupropion SR for smoking cessation in smokers with cardiovascular disease: a multicentre, randomised study. *Eur Heart J* 2003;24:946-55.

Transdermal Nicotine Study Group. Transdermal nicotine for smoking cessation. *Six*

Table 7.1. Evidence of effectiveness of tobacco dependence interventions in specific populations. Section 7, Psychiatric disorders including substance use disorders. (n = 4 studies)

Hall SM, Reus VI, Munoz RF, et al. Nortriptyline and cognitive-behavioral therapy in the treatment of cigarette smoking. Arch Gen Psychiatry 1998;55:683-90.

Hayford KE, Patten CA, Rummans TA, et al. Efficacy of bupropion for smoking cessation in smokers with a former history of major depression or alcoholism. Br J Psychiatry 1999;174:173-8.

Smith SS, Jorenby DE, Leischow SJ, et al. Targeting smokers at increased risk for relapse: treating women and those with a history of depression. Nicotine Tob Res 2003;5(1):99-109.

Wagena, EJ, Knipschild PG, Huibers MJ, et al. Efficacy of bupropion and nortriptyline for smoking cessation among people at risk for or with chronic obsth cnMh ce

