General References

1. Fiore MC, Bailey WC, Cohen SJ, et al. Smoking cessation. Clinical Practice Guideline No. 18. Rockville, MD, U.S. Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research. AHCPR Publication No. 96-0692. April 1996.
2. National Center for Health Statistics. Health, United States, 2006, with chartbook on trends in the health of Americans. Hyattsville, MD, 2006.
3. Centers for Disease Control and Prevention. Cigarette smoking among adults—United 2008;6:60-8.
6. U.S. Department of Health and Human Services. Reducing tobacco use: a report of the Surgeon General. Atlanta, GA, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.
7. Baumeister SE, Schumann A, Meyer C, et al. Effects of smoking cessation on health care use: is elevated risk of hospitalization among former smokers attributable to smoking-related morbidity? Drug Alcohol Depend 2007; 88:197-203.
8. Lightwood J. The economics of smoking and cardiovascular disease. Prog Cardiovasc Dis 2003;46:39-78.
9. Lightwood JM, Glantz SA. Short-term economic and health benefits of smoking cessation: myocardial infarction and stroke. Circulation 1997; 96:1089-96.
10. Rasmussen SR, Prescott E, Sorensen TI, et al. The total lifetime health cost savings of smoking cessation to society. Eur J Public Health 2005;15:601-6.
12. Critchley J, Capewell S. Smoking cessation for the secondary prevention of coronary heart disease. Cochrane Database Syst Rev 2004:CD003041.
13. Ford ES, Ajani UA, Croft JB, et al. Explaining the decrease in U.S. deaths from coronary disease, 1980-2000. N Engl J Med 2007, 356:2388-98.
14. Centers for Disease Control and Prevention. Cigarette smoking among adults—United States, 2004. MMWR 2005;54:1121-24.
15. Centers for Disease Control and Prevention. Cigarette smoking among adults—United States, 2003. MMWR 2005;54:509-13.
16. Tobacco Cessation Leadership Network. Trends in the delivery and reimbursement of tobacco dependence treatment. Available at: http://tcln.org/resources/pdfs/Trends_in_Delivery_and_Reimbursement_final.pdf.)
17. Centers for Disease Control and Prevention. State Medicaid coverage for tobacco-dependence treatments—United States, 2005. MMWR 2006;55:1194-7.
18. Bellows NM, McMenamin SB, Halpin HA. Adoption of system strategies for tobacco cessation by state Medicaid programs. Med Care 2007;45:350-6.
19. Centers for Disease Control and Prevention. Physician and other health care professional counseling of smokers to quit—United States, 1991. MMWR 1993;42:854-7.
20. Centers for Disease Control and Prevention. Receipt of advice to quit smoking in Medicare managed care—United States, 1998. JAMA 2000;284:1779-81.
21. Denny CH, Serdula MK, Holtzman D, et al. Physician advice about smoking and drinking: are U.S. adults being informed? Am J Prev Med 2003;24:71-4.
22. Chase EC, McMenamin SB, Halpin HA. Medicaid provider delivery of the 5A's for smoking cessation counseling. Nicotine Tob Res 2007,9;1095-101.
23. Quinn VP, Stevens VJ, Hollis JF, et al. Tobacco-cessation services and patient satisfaction in nine nonprofit HMOs. Am J Prev Med 2005;29:77-84.
24. California Department of Health Services. Smokers and quitting. 2005. Available at:

48. Thorndike AN, Rigotti NA, Stafford RS, et al. National patterns in the treatment of smokers by physicians. JAMA 1998;279:604-8.
49. Thorndike AN, Regan S, Rigotti NA. The treatment of smoking by US physicians during ambulatory visits: 1994 2003. Am J Public Health 2007;97:1878-83.
50. Conroy MB, Majchrzak NE, Regan S, et al. The association between patient-reported receipt of tobacco intervention at a primary care visit and smokers' satisfaction with their health care. Nicotine Tob Res 2005;7 Suppl 1:S29-34.
51. Stevens VJ, Solberg LI, Quinn VP, et al. Relationship between tobacco control policies and the delivery of smoking cessation services in nonprofit HMOs. J Natl Cancer Inst Monogr 2005:75-80.
52. Fang CY, Ma GX, Miller SM, et al. A brief smoking cessation intervention for Chinese and Korean American smokers. Prev Med 2006;43:321-4.
53. Ferreira-Borges C. Effectiveness of a brief counseling and behavioral intervention for smoking cessation in pregnant women. Prev Med 2005;41:295-302.
54. Fiore MC, Jorenby DE, Schensky AE, et al. Smoking status as the new vital sign: effect on assessment and intervention in patients who smoke. Mayo Clin Proc 1995;70:209-13.
55. Glasgow RE, Whitlock EP, Eakin EG, el al. A brief smoking cessation intervention for women in low-income planned parenthood clinics. Am J Public Health 2000;90:786-9.
56. Lancaster T, Stead L. Physician advice for smoking cessation. Cochrane Database Syst Rev 2004:CD000165.
57. Molyneux A, Lewis S, Leivers U, et al. Clinical trial comparing nicotine replacement therapy (NRT) plus brief counseling, brief counseling alone, and minimal intervention on smoking cessation in hospital inpatients. Thorax 2003;58:484-8.
58. Rissel C, Salmon A, Hughes AM. Evaluation of a (pilot) stage-tailored brief smoking cessation intervention among hospital patients presenting to a hospital pre-admission clinic. Aust Health Rev 2000;23:83-93.
59. Bonollo DP, Zapka JG, Stoddard AM, et al. Treating nicotine dependence during pregnancy and postpartum: understanding clinician knowledge and performance. Patient Edu Couns 2002;48:265-74.

100. Garvey AJ, Bliss RE, Hitchcock JL, et al. Predictors of smoking relapse among self-quitters: a report from the Normative Aging Study. Addict Behav 1992;17:367-77.
101. Westman EC, Behm FM, Simel DL, et al. Smoking behavior on the first day of a quit attempt predicts long-term abstinence. Arch Intern Med 1997;157:333-40.
102. Hurt RD, Krook JE, Croghan IT, et al. Nicotine patch therapy based on smoking rate followed by bupropion for prevention of relapse to smoking. J Clin Oncol 2003;21:914-20.
103. DerSimonian R, Laird N. Meta-analysis in clinical trials. Control Clin Trials 1986;7:177-88.
104. Statistics and Epidemiology Research Corporation. EGRET Reference Manual, Revision 4. Seattle, WA, 1993.
105. Hosmer DW, Lemeshow S. Applied Logistic Regression, Second Edition. New York, John Wiley & Sons, Inc., 2000.
106. Gartlehner G, Hanson RA, Nissman D, et al. Criteria for distinguishing effectiveness from efficacy trials is systematic reviews. Technical Review 12 (Prepared by the RTI-International-University of North Carolina Evidence-based Practice Center under Contract No. 290-02-0016.) AHRQ Publication No. 06-0046. Rockville, MD, Agency for Healthcare Research and Quality, April 2006.
107. Dale LC, Hurt RD, Hays JT. Drug therapy to aid in smoking cessation. Tips on maximizing patients' chances for success. Postgrad Med 1998;104:75-8, 83-74.
108. Lerman C, Kaufmann V, Rukstalis M, et al. Individualizing nicotine replacement therapy for the treatment of tobacco dependence: a randomized trial. Ann Intern Med 2004;140:426-33.
109. West R, Hajek P, Nilsson F, et al. Individual differences in preferences for and responses to four nicotine replacement products. Psychopharmacology (Berl) 2001;153:225-30.
110. Drilea SK, Reid BC, Li CH, et al. Dental visits among smoking and nonsmoking US adults in 2000. Am J Health Behav 2005;29:462-71.
111. Centers for Disease Control and Prevention. Cigarette smoking among adults—United States, 2000. MMWR 2002;51:642-5.
112. Fu SS, Partin MR, Snyder A, et al. Promoting repeat tobacco dependence treatment: are relapsed smokers interested? Am J Manag Care 2006;12:235-43.
113. National Cancer Institute. Tobacco and the clinician: interventions for medical and dental practice. Monograph No. 5. NIH Publication No. 94-3696, 1994.
114. Kreuter MW, Chheda SG, Bull FC. How does physician advice influence patient behavior? Evidence for a priming effect. Arch Fam Med 2000;9:426-33.
115. Richmond RL. Physicians can make a difference with smokers: evidence-based clinical approaches. Presentation given during the Symposium on Smoking Cessation at the 29th World Conference of the IUATLD/UICTMR and Global Congress on Lung Health, Bangkok, Thailand, 23-26 November 1998. International Union Against Tuberculosis and Lung Disease. Int J Tuberc Lung Dis 1999;3:100-12.
116. Whitlock EP, Orleans CT, Pender N, et al. Evaluating primary care behavioral counseling interventions: an evidence-based approach. Am J Prev Med 2002;22:267-84.
117. Tessaro I, Lyna PR, Rimer BK, et al. Readiness to change smoking behavior in a community health center population. J Community Health 1997;22:15-31.
118. Ossip-Klein DJ, McIntosh S, Utman C, et al. Smokers ages 50+: who gets physician advice to quit? Prev Med 2000;31:364-9.
119. National Committee for Quality Assurance. The state of health care quality 2006. Available at: http://www.ncqa.org/communications/sohc2006/sohc_2006.pdf.
120. Shelley D, Cantrell J, Faulkner D, et al. Physician and dentist tobacco use counseling and adolescent smoking behavior: results from the 2000 National Youth Tobacco Survey. Pediatrics 2005;115:719-25.
121. Moran S, Thorndike AN, Armstrong K, et al. Physicians' missed opportunities to address tobacco use during prenatal care. Nicotine Tob Res 2003;5:363-8.
122. Rennard SI, Daughton DM. Smoking cessation. Chest 2000;117:360S-4S.
123. Doll R, Peto R, Boreham J, et al. Mortality in relation to smoking: 50 years' observations on male British doctors. BMJ 2004;328:1519.
124. Lichtenstein E, Hollis J. Patient referral to a smoking cessation program: who follows through? J Fam Prac 1992;34:739-44.
125. Hammond D, McDonald PW, Fong GT, et al. Do smokers know how to quit? Knowledge and perceived effectiveness of cessation assistance as predictors of cessation behaviour. Addiction 2004;99:1042-8.

126. Roche AM, Freeman T. Brief interventions: good in theory but weak in practice. Drug Alcohol Rev 2004;23:11-8.
127. Stange KC, Zyzanski SJ, Jaen CR, et al. Illuminating the ‘black box’. A description of 4,454 patient visits to 138 family physicians. J Fam Pract 1998;46:377-89.
128. Nides M, Leischow S, Sarna L, et al. Maximizing smoking cessation in clinical practice: pharmacologic and behavioral interventions. Prev Cardiol 2007;10:23-30.
129. Tomar SL, Husten CG, Manley MW. Do dentists and physicians advise tobacco users to quit? J Am Dent Assoc 1996;127:259-65.
130. Schroeder SA. What to do with a patient who smokes. JAMA 2005;294:482-7.
131. Gordon JS, Andrews JA, Crews KM, et al. The 5A’s vs 3A’s plus proactive quitline referral in private practice dental offices: preliminary results. Tob Control 2007;16:285-8.
patients stop smoking: a National Cancer Institute manual for physicians. Bethesda, MD, NIH Publication No. 89-3064, 1997.
133tl(pID 4 Manley MW,)6(Epps RP, BD)9(nn TJ. The)]TJ0.0005 Tc -0.0029 Tw 2.695 -1.15 Td[(clini)7(c)-2(ian’)7(s ro

180. Carpenter MJ, Hughes JR, Keely JP. Effect of smoking reduction on later cessation: a pilot experimental study. Nicotine Tob Res 2003;5:155-62.
181. Piasecki TM, Fiore MC, McCarthy DE, et al. Have we lost our way? The need for dynamic formulations of smoking relapse proneness. Addiction 2002;97:1093-108.
182. Haddock CK, O'Byrne KK, Klesges RC, et al. Relapse to smoking after basic military training in the U.S. Air Force. Mil Med 2000;165:884-8.
183. Krall EA, Garvey AJ, Garcia RI. Smoking relapse after 2 years of abstinence: findings from the VA Normative Aging Study. Nicotine Tob Res 2002;4:95-100.
184. Wetter DW, Cofta-Gunn L, Fouladi RT, et al. Late relapse/sustained abstinence among former smokers: a longitudinal study. Prev Med 2004;39:1156-63.
185. Hajek P, Stead LF, West R, et al. Relapse prevention interventions for smoking cessation. Cochrane Database Syst Rev 2005:CD003999
186. Brandon TH, Collins BN, Juliano LM, et al. Preventing relapse among former smokers: a comparison of minimal interventions through telephone and mail. J Consult Clin Psychol 2000;68:103-13.
187. Brandon TH, Meade CD, Herzog TA, et al. Efficacy and cost-effectiveness of a minimal intervention to prevent smoking relapse: dismantling the effects of amount of content versus contact. J Consult Clin Psychol 2004;72:797-808.
188. Alterman AI, Gariti P, Mulvaney F. Short- and long-term smoking cessation for three levels of intensity of behavioral treatment. Psychol Addict Behav 2001;15:261-4.
189. Barth J, Critchley J, Bengel J. Efficacy of psychosocial interventions for smoking cessation in patients with coronary heart disease: a systematic review and meta-analysis. Ann Behav Med 2006;32:10-20.
190. Grandes G, Cortada JM, Arrazola A, et al. Predictors of long-term outcome of a smoking cessation programme in primary care. Br J Gen Pract 2003;53:101-7.
191. Hilleman DE, Mohiuddin SM, Packard KA. Comparison of conservative and aggressive smoking cessation treatment strategies following coronary artery bypass graft surgery. Chest 2004;125:435-8.
192. Pisinger C, Vestbo J, Borch-Johnsen K, et al. Smoking cessation intervention in a large randomised population-based study. The Inter99 study. Prev Med 2005;40:285-92.
193. Rigotti NA, Munafo MR, Murphy MF, et al. Interventions for smoking cessation in hospitalized patients. Cochrane Database Syst Rev 2003;1:1-21.
194. Cromwell J, Bartosch WJ, Fiore MC, et al. Cost-effectiveness of the clinical practice recommendations in the AHCPR guideline for smoking cessation. JAMA 1997;278:1759-66.
195. Bentz CJ, Bayley KB, Bonin KE, et al. The feasibility of connecting physician offices to a state-level tobacco quit line. Am J Prev Med 2006;30:31-7.
196. Holtrop JS, Wadland WC, Vansen S, et al. Recruiting health plan members receiving pharmacotherapy into smoking cessation counseling. Am J Manag Care 2005;11:501-7.
197. Perry RJ, Keller PA, Fraser D, et al. Fax to quit: a model for delivery of tobacco cessation services to Wisconsin residents. WMJ 2005;104:37-40, 44.
198. Winickoff JP, Buckley VJ, Palfrey JS, et al. Intervention with parental smokers in an outpatient pediatric clinic using counseling and nicotine replacement. Pediatrics 2003;112:1127-33.
199. Winickoff JP, Hillis VJ, Palfrey JS, et al. A smoking cessation intervention for parents of children who are hospitalized for respiratory illness: The Stop Tobacco Outreach Program.

205. Levy DT, Chaloupka F, Gitchell J. The effects of tobacco control policies on smoking rates: a tobacco control scorecard. J Public Health Manag Pract 2004;10:338-53.
206. Curry SJ, Orleans CT, Keller P, et al. Promoting smoking cessation in the healthcare environment: 10 years later. Am J Prev Med 2006;31:269-72.
207. Zapka JG, White MJ, Reed G, et al. Organizational systems to support publicly funded tobacco treatment services. Am J Prev Med 2005;28:338-45.
208. Wewers ME, Kidd K, Armbruster D, et al. Tobacco dependence curricula in U.S. baccalaureate and graduate nursing education. Nurs Outlook 2004;52:95-101.
209. Glasgow RE, Davis CL, Funnell MM, et al. Implementing practical interventions to support chronic illness self-management. Jt Comm J Qual Saf 2003;29:563-74.
210. Curry SJ, Grothaus LC, McAfee T, et al. Use and cost effectiveness of smoking-cessation services under four insurance plans in a health maintenance organization. N Engl J Med 1998;339:673-9.
211. Doescher MP, Whinston MA, Goo A, et al. Pilot study of enhanced tobacco-cessation services coverage for low-income smokers. Nicotine Tob Res 2002;4 Suppl 1:S19-24.
212. Kaper J, Wagena EJ, Severens JL, et al. Health care financing systems for increasing the use of tobacco dependence treatment. Cochrane Database Syst Rev 2005:CD004305.
213. Manley MW, Griffin T, Foldes SS, et al. The role of health plans in tobacco control. Annu Rev Public Health 2003;24:247-66.
214. McPhillips-Tangum C, Rehm B, Carreon R, et al. Addressing tobacco in managed care: results of the 2003 survey. Prev Chronic Dis 2006;3:A87.
215. Bondi MA, Harris JR, Atkins D, et al. Employer coverage of clinical preventive services in the United States. Am J Health Promot 2006;20:214-22.
216. Halpin HA, Bellows NM, McMenamin SB. Medicaid coverage for tobacco-dependence treatments. Health Aff (Millwood) 2006;25:550-6.
217. Centers for Disease Control and Prevention. State Medicaid coverage for tobacco-dependence treatments—United States, 1994-2002. MMWR 2004;53:54-7.
218. Ibrahim JK, Schauffler HH, Barker DC, et al. Coverage of tobacco dependence treatments for pregnant women and for children and their parents. Am J Public Health 2002;92:1940-2.
219. Centers for Disease Control and Prevention. State Medicaid coverage for tobacco-dependence treatments—United States, 2005. MMWR 2006;55:1194-7.
220. U.S. Department of Health and Human

the Pacific Business Group on Health. Am J Health Promot 2001;15:350-6.
232. Ahluwalia JS, Gibson CA, Kenney RE, et al. Smoking status as a vital sign. J Gen Intern Med 1999;14:402-8.
233. Boyle R, Solberg LI. Is making smoking status a vital sign sufficient to increase cessation support actions in clinical practice? Ann Fam Med 2004;2:22-5.
234. Glasgow RE, Lichtenstein E, Marcus AC. Why don't we see more translation of health promotion research to practice? Rethinking the efficacy-to-effectiveness transition. Am J Public Health 2003;93:1261-7.
235. Glasgow RE, Orleans CT, Wagner EH. Does the chronic care model serve also as a template for improving prevention? Milbank Q 2001;79:579-612, iv-v.
236. Keller PA, Fiore MC, Curry SJ, et al. Systems change to improve health and health care: lessons from addressing tobacco in managed care. Nicotine Tob Res 2005;7 Suppl 1:S5-8.
237. Piper ME, Fiore MC, Smith SS, et al. Use of the vital sign stamp as a systematic screening tool to promote smoking cessation. Mayo Clin Proc 2003;78:716-22.
238. Borland R, Balmford J, Hunt D. The effectiveness of personally tailored computer-generated advice letters for smoking cessation. Addiction 2004;99:369-77.
239. Kim JR, Lee MS, Hwang JY, et al. Efficacy of a smoking cessation intervention using the AHCPR guideline tailored for Koreans: a randomized controlled trial. Health Promot Int 2005;20:51-9.
240. McClure JB, Swan GE. Tailoring nicotine replacement therapy: rationale and potential approaches. CNS Drugs 2006;20:281-91.
241. Strecher VJ, Marcus A, Bishop K, et al. A randomized controlled trial of multiple tailored messages for smoking cessation among callers to the cancer information service. J Health Commun 2005;10 Suppl 1:105-18.
242. Strecher VJ, Shiffman S, West R. Moderators and mediators of a web-based computer-tailored smoking cessation program among nicotine patch users. Nicotine Tob Res 2006;8 Suppl 1:S95-101.
243. Unrod M, Smith M, Spring B, et al. Randomized controlled trial of a computer-based, tailored intervention to increase smoking cessation counseling by primary care physicians. J Gen Intern Med 2007;22:478-84.
244. Webb MS, Simmons VN, Brandon TH. Tailored interventions for motivating smoking cessation: using placebo tailoring to examine the influence of expectancies and personalization. Health Psychol 2005;24:179-88.
245. Velicer WF, Prochaska JO, Redding CA. Tailored communications for smoking cessation: past successes and future directions. Drug Alcohol Rev 2006;25:49-57.
246. Gilbert DG, Crauthers DM, Mooney DK, et al. Effects of monetary contingencies on smoking relapse: influences of trait depression, personality, and habitual nicotine intake. Exp Clin Psychopharmacol 1999;7:174-81.
247. Han ES, Foulds J, Steinberg MB, et al. Characteristics and smoking cessation outcomes of patients returning for repeat tobacco dependence treatment. Int J Clin Pract 2006;60:1068-74.
248. Smith SS, Jorenby DE, Leischow SJ, et al. Targeting smokers at increased risk for relapse: treating women and those with a history of depression. Nicotine Tob Res 2003;5:99-109.
249. Bao Y, Duan N, Fox SA. Is some provider advice on smoking cessation better than no advice? An instrumental variable analysis of the 2001 National Health Interview Survey. Health Serv Res 2006;41:2114Tɔ Tw c5TJEMC /P <</MCII

257. Smith SS, Jorenby DE, Fiore MC, et al. Strike while the iron is hot: can stepped-care treatments resurrect relapsing smokers? J Consult Clin Psychol 2001;69:429-39.
258. Reid R, Pipe A, Higginson L, et al. Stepped care approach to smoking cessation in patients hospitalized for coronary artery disease. J Cardiopulm Rehabil 2003;23:176-82.
259. Walters ST, Wright JA, Shegog R. A review of computer and Internet-based interventions for smoking behavior. Addict Behav 2006;31:264-77.
260. Lenert L, Munoz RF, Perez JE, et al. Automated e-mail messaging as a tool for improving quit rates in an internet smoking cessation intervention. J Am Med Inform Assoc 2004;11:235-40.
261. Bock B, Graham A, Sciamanna C, et al. Smoking cessation treatment on the Internet: content, quality, and usability. Nicotine Tob Res 2004;6:207-19.
262. Etter JF. Comparing the efficacy of two Internet-based, computer-tailored smoking cessation programs: a randomized trial. J Med Internet Res 2005;7:e2
263. Cobb NK, Graham AL, Bock BC, et al. Initial evaluation of a real-world Internet smoking cessation system. Nicotine Tob Res 2005;7:207-16.
264. Japuntich SJ, Zehner ME, Smith SS, et al. Smoking cessation via the Internet: a randomized clinical trial of an internet intervention as adjuvant treatment in a smoking cessation intervention. Nicotine Tob Res 2006;8 Suppl 1:S59-67.
265. Graham AL, Bock BC, Cobb NK, et al. Characteristics of smokers reached and recruited to an Internet smoking cessation trial: a case of denominators. Nicotine Tob Res 2006;8 Suppl 1:S43-8.
266. Hall SM, Tsoh JY, Prochaska JJ, et al. Treatment for cigarette smoking among depressed mental health outpatients: a randomized clinical trial. Am J Public Health 2006;96:1808-14.
267. Prochaska JO, Velicer WF, Fava JL, et al. Counselor and stimulus control enhancements of a stage-matched expert system intervention for smokers in a managed care setting. Prev Med 2001;32:23-32.
268. Swartz SH, Cowan TM, Batista IA. Using claims data to examine patients using practice-based Internet communication: is there a clinical digital divide? J Med Internet Res 2004;6:e1.
269. Hollis JF, Polen MR, Whitlock EP, et al. Teen reach: outcomes from a randomized,

282. Abbot NC, Stead LF, White AR, et al. Hypnotherapy for smoking cessation. The

310. Johnstone E, Benowitz N, Cargill A, et al. Determinants of the rate of nicotine metabolism and effects on smoking behavior. Clin Pharmacol Ther 2006;80:319-30.

362. Zhu S, Melcer T, Sun J, Rosbrook B, Pierce

complementary health intervention: the case of smoking relapse prevention. Int J Technol Assess Health Care 2004;20:475-80.
390. Halpin HA, McMenamin SB, Rideout J, et al. The costs and effectiveness of different benefit designs for treating tobacco dependence: results from a randomized trial. Inquiry 2006;43:54-65.
391. Schauffler HH, McMenamin S, Olson K, et al. Variations in treatment benefits influence smoking cessation: results of a randomised controlled trial. Tob Control 2001;10:175-80.
392. Fung PR, Snape-Jenkinson SL, Godfrey MT, et al. Effectiveness of hospital-based smoking cessation. Chest 2005;128:216-23.
393. Lennox AS, Osman LM, Reiter E, et al. Cost

smoking cessation? Health Aff (Millwood) 2002;21:162-8.
417. An LC, Schillo BA, Kavanaugh AM, et al. Increased reach and effectiveness of a statewide tobacco quitline after the addition of access to free nicotine replacement therapy. Tob Control 2006;15:286-93.
418. Vidrine DJ, Arduino RC, Gritz ER. Impact of a cell phone intervention on mediating mechanisms of smoking cessation in individuals living with HIV/AIDS. Nicotine Tob Res 2006;8 Suppl 1:S103-8.
419. Elzi L, Spoerl D, Voggensperger J, et al. A smoking cessation programme in HIV-infected individuals: a pilot study. Antivir Ther 2006;11:787-95.
420. Rigotti N, Munafo M, Stead L. Interventions for smoking cessation in hospitalised patients. Cochrane Database Syst Rev 2007.
421. Taylor CB, Miller NH, Herman S, et al. A nurse-managed smoking cessation program for hospitalized smokers. Am J Public Health 1996;86:1557-60.
422. Ratner PA, Johnson JL, Richardson CG, et al. Efficacy of a smoking-cessation intervention for elective-surgical patients. Res Nurs Health 2004;27:148-61.
423. Simon JA, Carmody TP, Hudes ES, et al. Intensive smoking cessation counseling versus minimal counseling among hospitalized smokers treated with transdermal nicotine replacement: a randomized trial. Am J Med 2003;114:555-62.
424. Stevens VJ, Glasgow RE, Hollis JF, et al. A smoking-cessation intervention for hospital patients. Med Care 1993;31:65-72.
425. Miller NH, Smith PM, DeBusk RF, et al. Smoking cessation in hospitalized patients. Results of a randomized trial. Arch Intern Med 1997;157:409-15.
426. Groner JA, Ahijevych K, Grossman LK, et al. The impact of a brief intervention on maternal smoking behavior. Pediatrics 2000;105:267-71.
427. Solomon LJ, Secker-Walker RH, Flynn BS, et al. Proactive telephone peer support to help pregnant women stop smoking. Tob Control 2000;9 Suppl 3:III72-4.
428. Solomon LJ, Marcy TW, Howe KD, et al. Does extended proactive telephone support increase smoking cessation among low-income women using nicotine patches? Prev Med 2005;40:306-13.
429. Davies SL, Kohler CL, Fish L, et al. Evaluation of an intervention for hospitalized African American smokers. Am J Health Behav 2005;29:228-39.
430. Ockene J, Kristeller JL, Goldberg R, et al. Smoking cessation and severity of disease: the Coronary Artery Smoking Intervention Study. Health Psychol 1992;11:119-26.
431. Taylor CB, Miller NH, Killen JD, et al. Smoking cessation after acute myocardial infarction: effects of a nurse-managed intervention. Ann Intern Med 1990;113:118-23.
432. DeBusk RF, Miller NH, Superko HR, et al. A case-management system for coronary risk factor modification after acute myocardial infarction. Ann Intern Med 1994;120:721-9.
433. Chouinard MC, Robichaud-Ekstrand S. The effectiveness of a nursing inpatient smoking cessation program in individuals with cardiovascular disease. Nurs Res 2005;54:243-54.
434. Dornelas EA, Sampson RA, Gray JF, et al. A randomized controlled trial of smoking cessation counseling after myocardial infarction. Prev Med 2000;30:261-8.
435. Sivarajan Froelicher ES, Miller NH, Christopherson DJ, et al. High rates of sustained smoking cessation in women hospitalized with cardiovascular disease: the Women's Initiative for Nonsmoking (WINS). Circulation 2004;109:587-93.
436. Quist-Paulsen P, Gallefoss F. Randomised controlled trial of smoking cessation intervention after admission for coronary heart disease. BMJ 2003, 327:1254-1257
437. Giallauria F, Paragliola T, Pilerci F, et al. Role of smokers in the household and of cardiac rehabilitation in smoking behaviour after acute myocardial infarction. Monaldi Arch Chest Dis 2005;64:110-5.
438. McHugh F, Lindsay GM, Hanlon P, et al. Nurse led shared care for patients on the waiting list for coronary artery bypass surgery: a randomised controlled trial. Heart 2001;86:317-23.
439. Ludvig J, Miner B, Eisenberg MJ. Smoking cessation in patients with coronary artery disease. Am Heart J 2005;149:565-72.
440. Taylor CB, Houston-Miller N, Haskell WL, et al. Smoking cessation after acute myocardial infarction: The effects of exercise training. Addict Behav 1988;13:331-5.
441. Fowler B, Jamrozik K, Norman P, et al. Improving maximum walking distance in early peripheral arterial disease: randomised controlled trial. Aust J Physiother 2002;48:269-75.

442. Tonstad S, Farsang C, Klaene G, et al. Bupropion SR for smoking cessation in smokers with cardiovascular disease: a multicentre, randomised study. Eur Heart J 2003;24:946-55.
443. Rigotti NA, Thorndike AN, Regan S, et al. Bupropion for smokers hospitalized with acute cardiovascular disease. Am J Med 2006;119:1080-7.
444. Christenhusz L, Pieterse M, Seydel E, et al. Prospective determinants of smoking cessation in COPD patients within a high intensity or a brief counseling intervention. Patient Educ Couns 2007;66:162-6.
445. Wagena EJ, van der Meer RM, Ostelo RJ, et al. The efficacy of smoking cessation strategies in people with chronic obstructive pulmonary disease: results from a systematic review. Respir Med 2004;98:805-15.
446. Tashkin D, Kanner R, Bailey W, et al. Smoking cessation in patients with chronic obstructive pulmonary disease: a double-blind, placebo-controlled, randomised trial. Lancet 2001;357:1571-5.
447. Wagena EJ, Knipschild PG, Huibers MJ, et al. Efficacy of bupropion and nortriptyline for smoking cessation among people at risk for or with chronic obstructive pulmonary disease. Arch Intern Med 2005;165:2286-92.
448. Tonnesen P, Mikkelsen KL. Smoking cessation with four nicotine replacement regimes in a lung clinic. Eur Respir J 2000;16:717-22.
449. Duffy SA, Ronis DL, Valenstein M, et al. A tailored smoking, alcohol, and depression intervention for head and neck cancer patients. Cancer Epidemiol Biomarkers Prev 2006;15:2203-8.
450. Park ER, Puls-ul007 Trutvenfie5(3o)u;cctive detpeer-ductilw Tre028 Tline for

467. Evins AE, Cather C, Rigotti NA, et al. Two-year follow-up of a smoking cessation trial in patients with schizophrenia: increased rates of smoking cessation and reduction. J Clin Psychiatry 2004;65:307-11; quiz 452-303.
468. Kisely SR, Wise M, Preston N, et al. A group intervention to reduce smoking in individuals with psychiatric disorder: brief report of a pilot study. Aust N Z J Public Health 2003;27:61-3.
469. Evins AE, Mays VK, Rigotti NA, et al. A pilot trial of bupropion added to cognitive behavioral therapy for smoking cessation in schizophrenia. Nicotine Tob Res 2001;3:397-403.
470. George TP, Vessicchio JC, Termine A, et al. A placebo controlled trial of bupropion for smoking cessation in schizophrenia. Biol Psychiatry 2002;52:53-61.
471. Chou KR, Chen R, Lee JF, Ku CH, Lu RB. The effectiveness of nicotine-patch therapy for smoking cessation in patients with schizophrenia. Int J Nurs Stud 2004;41:321-30.
472. George TP, Ziedonis DM, Feingold A, et al. Nicotine transdermal patch and atypical antipsychotic medications for smoking cessation in schizophrenia. Am J Psychiatry 2000;157:1835-42.
473. Orleans CT, Hutchinson D. Tailoring nicotine addiction treatments for chemical dependency patients. J Subst Abuse Treat 1993;10:197-208.
474. Burling TA, Burling AS, Latini D. A controlled smoking cessation trial for substance-dependent inpatients. J Consult Clin Psychol 2001;69:295-304.
475. Hurt RD, Eberman KM, Croghan IT, et al. Nicotine dependence treatment during inpatient treatment for other addictions: a prospective intervention trial. Alcohol Clin Exp Res 1994;18:867-72.
476. Sussman S. Smoking cessation among persons in recovery. Subst Use Misuse 2002;37:1275-98.
477. Bobo JK, McIlvain HE, Lando HA, et al. Effect of smoking cessation counseling on recovery from alcoholism: findings from a randomized community intervention trial. Addiction 1998;93:877-87.
478. Ellingstad TP, Sobell LC, Sobell MB, et al. Alcohol abusers who want to quit smoking: implications for clinical treatment. Drug Alcohol Depen 1999;54:259-65.

cessation: a randomized controlled trial in Spanish smokers. Addict Behav 2002;27:139-44.
493. Munoz RF, Marin BV, Posner SF, et al. Mood management mail intervention increases abstinence rates for Spanish-speaking Latino smokers. Am J Community Psychol 1997;25:325-43.
494. Woodruff SI, Talavera GA, Elder JP. Evaluation of a culturally appropriate smoking cessation intervention for Latinos. Tob Control 2002;11:361-8.
495. Johnson KM, Lando HA, Schmid LS, Solberg LI: The GAINS project: outcome of smoking cessation strategies in four urban Native American clinics. Giving American Indians no-smoking strategies. Addictive Behaviors 1997, 22:207-218
496. Gonzales D, Rennard SI, Nides M, et al. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs sustained-release bupropion and placebo for smoking cessation: a randomized controlled trial. JAMA 2006;296:47-55.
497. Scharf D, Shiffman S. Are there gender differences in smoking cessation, with and without bupropion? Pooled- and meta-analyses of clinical trials of Bupropion SR. Addiction 2004;99:1462-9.
498. Wetter DW, Kenford SL, Smith SS, et al. Gender differences in smoking cessation. J Consult Clin Psychol 1999;67:555-62.
499. Copeland AL, Martin PD, Geiselman PJ, et al. Smoking cessation for weight-concerned women: group vs. individually tailored, dietary, and weight-control follow-up sessions. Addict Behav 2006;31:115-27.
500. Yilmaz G, Karacan C, Yoney A, et al. Brief intervention on maternal smoking: a randomized controlled trial. Child Care Health Dev 2006;32:73-9.
501. Marcus BH, Albrecht AE, King TK, et al. The efficacy of exercise as an aid for smoking cessation in women: a randomized controlled trial. Arch Intern Med 1999;159:1229-34.
502. Marcus BH, Albrecht AE, Niaura RS, et al. Exercise enhances the maintenance of smoking cessation in women. Addict Behav 1995;20:87-92.
503. Keller PA, Bailey LA, Koss KJ, et al. Organization, financing, promotion, and cost of U.S. quitlines, 2004. Am J Prev Med 2007;32:32-7.
504. Pierce J, Giovino G, Hatziandreu E, et al. National age and sex differences in quitting smoking. Journal of Psychoactive Drugs 1989;21:293-8.
505. Sorensen G, Barbeau E, Hunt MK, et al. Reducing social disparities in tobacco use: a social-contextual model for reducing tobacco use among blue-collar workers. Am J Public Health 2004;94:230-9.
506. Doolan DM, Froelicher ES. Efficacy of smoking cessation intervention among special populations: review of the literature from 2000 to 2005. Nurs Res 2006;55:S29-37.
507. Burkhalter JE, Springer CM, Chhabra R, et al. Tobacco use and readiness to quit smoking in low-income HIV-infected persons. Nicotine Tob Res 2005;7:511-22.
508. Mamary EM, Bahrs D, Martinez S. Cigarette smoking and the desire to quit among individuals living with HIV. Aids Patient Care STDS 2002;16:39-42.
509. Niaura R, Shadel WG, Morrow K, et al. Human immunodeficiency virus infection, AIDS, and smoking cessation: the time is now. Clin Infect Dis 2000;31:808-12.
510. Gritz ER, Vidrine DJ, Lazev AB, et al. Smoking behavior in a low-income multiethnic HIV/AIDS population. Nicotine Tob Res 2004;6:71-7.
511. Boerma JT, Nunn AJ, Whitworth JA. Mortality impact of the AIDS epidemic: evidence from community studies in less developed countries. AIDS 1998;12 Suppl 1:S3-14.
Reference Manager 11 (Build 6.0.1.1443(Dool)6
5(52[(sa(CM,Talavera,A,Reb(ter-VB,et al)8()5()a(R,
ceir tedu fmeb gn i kon
S

517. Hajjeh RA, Conn LA, Stephens DS, et al. Cryptococcosis: population-based multistate active surveillance and risk factors in human immunodeficiency virus-infected persons. Cryptococcal Active Surveillance Group. J Infect Dis 1999;179:449-54.
518. Shiboski CH, Neuhaus JM, Greenspan D, et al. Effect of receptive oral sex and smoking on the incidence of hairy leukoplakia in HIV-positive gay men. J Acquir Immune Defic Syndr 1999;21:236-42.
519. Chattopadhyay A, Caplan DJ, Slade GD, et al. Incidence of oral candidiasis and oral hairy leukoplakia in HIV-infected adults in North Carolina. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2005;99:39-47.
520. Slavinsky J 3rd, Myers T, Swoboda RK, et al. Th1/Th2 cytokine profiles in saliva of HIV-positive smokers with oropharyngeal candidiasis. Oral Microbiol Immunol 2002;17:38-43.
521. Metersky ML, Colt HG, Olson LK, et al. AIDS-related spontaneous pneumothorax. Risk factors and treatment. Chest 1995;108:946-51.
522. Reynolds NR, Neidig JL, Wewers ME. Illness representation and smoking behavior: a focus group study of HIV-positive men. J Assoc Nurses AIDS Care 2004;15:37-47.
523. Suskin N, Sheth T, Negassa A, et al. Relationship of current and past smoking to mortality and morbidity in patients with left ventricular dysfunction. J Am Coll Cardiol 2001;37:1677-82.
524. Reid ME, Marshall JR, Roe D, et al. Smoking exposure as a risk factor for prevalent and recurrent colorectal adenomas. Cancer Epidemiol Biomarkers Prev 2003;12:1006-11.
525. Ryan WR, Ley C, Allan RN, et al. Patients with Crohn's disease are unaware of the risks that smoking has on their disease. J Gastrointest Surg 2003;7:706-11.
526. Pacchioni D, Martone T, Ghisolfi G, et al. Tobacco smoke, recurrences, and p53/bcl-2 expression in bladder cancer. Carcinogenesis 1997;18:1659-61.
527. Browman GP, Mohide EA, Willan A, et al. Association between smoking during radiotherapy and prognosis in head and neck cancer: a follow-up study. Head Neck 2002;24:1031-7.
528. Gritz ER, Carr CR, Rapkin D, et al. Predictors of long-term smoking cessation in head and neck cancer patients. Cancer Epidemiol Biomarkers Prev 1993;2:261-70.
529. Kawahara M, Ushijima S, Kamimori T, et al. Second primary tumours in more than 2-year disease-free survivors of small-cell lung cancer in Japan: the role of smoking cessation. Br J Cancer 1998;78:409-12.
530. Richardson GE, Tucker MA, Venzon DJ, et al. Smoking cessation after successful treatment of small-cell lung cancer is associated with fewer smoking-related second primary cancers. Ann Intern Med 1993;119:383-90.
531. Rennard SI, Togo S, Holz O. Cigarette smoke inhibits alveolar repair: a mechanism for the development of emphysema. Proc Am Thorac Soc 2006;3:703-8.
532. Ziran BH, Hendi P, Smith WR, et al. Osseous healing with a composite of allograft and demineralized bone matrix: adverse effects of smoking. Am J Orthop 2007;36:207-9.
533. Moller A, Tonnesen H. Risk reduction: perioperative smoking intervention. Best Pract Res Clin Anaesthesiol 2006;20:237-48.
534. Rogliani M, Labardi L, Silvi E, et al. Smokers: risks and complications in abdominal dermolipectomy. Aesthetic Plast Surg 2006;30:422-4; discussion 425.
535. Levin L, Schwartz-Arad D. The effect of cigarette smoking on dental implants and related surgery. Implant Dent 2005;14:357-61.
536. Spear SL, Ducic I, Cuoco F, et al. The effect of smoking on flap and donor-site complications in pedicled TRAM breast reconstruction. Plast Reconstr Surg 2005;116:1873-80.
537. Willemse BW, Ten Hacken NH, Rutgers B, et al. Effect of 1-year smoking cessation on airway inflammation in COPD and asymptomatic smokers. Eur Respir J 2005;26:835-45.
538. Scanlon PD, Connett JE, Waller LA, et al. Smoking cessation and lung function in mild-to-moderate chronic obstructive pulmonary disease. The Lung Health Study. Am J Respir Crit Care Med 2000;161:381-90.
539. Emmons KM, Butterfield RM, Puleo E, et al. Smoking among participants in the childhood cancer survivors cohort: the Partnership for Health Study. J Clin Oncol 2003;21:189-96.
540. Hurt RD, Berge KG, Offord KP, et al. The making of a smoke-free medical center. JAMA 1989;261:95-7.
541. Maryland Hospital Association. Smoke-free hospital campus tool-kit. Available at: http://www.mdhospitals.org/mha/Community_

Health_Resources/Smoke_Free_Hospital_Campuses.shtml.
542. Michigan Health and Hospital Association. It's a matter of life & health: MHA campaign for smoke-free hospitals. Available at: http://www.mha.org/mha_app/public_site/smokefree.jsp.
543. Suplee PD. The importance of providing smoking relapse counseling during the

567. Hoffman C, Carbaugh A, Cook A. Health insurance coverage in America: 2003 data update. Kaiser Commision on Medicaid and the Unisured. Kaiser Commision on Medicaid and the Unisured. Washington, DC, Kaiser Family Foundation, 2004
568. Wadland WC, Soffelmayr B, Ives K. Enhancing smoking cessation of low-income smokers in managed care. J Fam Prac 2001;50:138-44.
569. Dermer A. Smoking, tobacco exposure through breast milk, and SIDS. JAMA 1995;274:214-5.
570. Jaen CR, Cummings KM, Shah D, et al. Patterns of use of a free nicotine patch program for Medicaid and uninsured patients. J Natl Med Assoc 1997;89:325-8.
571. Leone A. Smoking, haemostatic factors, and cardiovascular risk. Curr Pharm Des 2007;13:1661-7.
572. Lavie L, Lavie P. Smoking interacts with sleep apnea to increase cardiovascular risk. Sleep Med 2007;May 18; [Epub ahead of print]
573. Wilson PW. Smoking, smoking cessation, and risk of cardiovascular disease. Curr Treat Options Cardiovasc Med 2006;8:276-81.
574. Twardella D, Rothenbacher D, Hahmann H, et al. The underestimated impact of smoking and smoking cessation on the risk of secondary cardiovascular disease events in patients with stable coronary heart disease:

Handbooks on Cancer Prevention No. 11, World Health Organization, 2007.
594. Kuri M, Nakagawa M, Tanaka H, et al. Determination of the duration of preoperative smoking cessation to improve wound healing after head and neck surgery. Anesthesiology 2005;102:892-6.
595. Yamashita K, Kobayashi S, Yamaguchi S. Cerebral blood flow and cessation of cigarette smoking in healthy volunteers. Intern Med 2000;39:891-3.
596. Hall SM, Humfleet GL, Gorecki JA, et al. Older versus younger treatment-seeking smokers: differences in smoking behavior, drug and alcohol use, and psychosocial and physical functioning. Nicotine Tob Res

648. Anderson C, Burns DM. Patterns of adolescent smoking initiation rates by ethnicity and sex. Tob Control 2000;9 Suppl 2:II4-8.
649. Barbeau EM, Krieger N, Soobader MJ. Working class matters: socioeconomic disadvantage, race/ethnicity, gender, and smoking in NHIS 2000. Am J Public Health 2004;94:269-78.
650. Watson JM, Scarinci IC, Klesges RC, et al. Relationships among smoking status, ethnicity, socioeconomic indicators, and lifestyle variables in a biracial sample of women. Prev Med 2003;37:138-47.
651. King G, Polednak A, Bendel RB, et al. Disparities in smoking cessation between African Americans and whites: 1990-2000. Am J Public Health 2004;94:1965-71.
652. Benowitz NL, Perez-Stable EJ, Herrera B, et al. Slower metabolism and reduced intake of nicotine from cigarette smoking in Chinese Americans. J Natl Cancer Inst 2002;94:108-15.
653. Lemogoum D, Van Bortel L, Leeman M, et al. Ethnic differences in arterial stiffness and wave reflections after cigarette smoking. J Hypertens 2006;24:683-9.
654. Lea R, Benowitz N, Green M, et al. Ethnic differences in nicotine metabolic rate among New Zealanders. N Z Med J 2005;118:U1773.
655. Nakajima M, Yokoi T. Interindividual variability in nicotine metabolism: C-oxidation and glucuronidation. Drug Metab Pharmacokinet 2005;20:227-35.
656. Mwenifumbo JC, Sellers EM, Tyndale RF. Nicotine metabolism and CYP2A6 activity in a population of black African descent: impact of gender and light smoking. Drug Alcohol Depend 2007;89:24-33.

population-based analysis. Am J Prev Med 2000;18:305-11.
674. Centers for Disease Control and Prevention. Youth and tobacco use: current estimates. 2006. Available at: http://www.cdc.gov/tobacco/youth/index.htm.
675. American Lung Association. Adolescent smoking statistics 2003 Available at: http://www.lungusa.org/site/apps/s/content.asp?c=dvLUK9O0E&b=34706&ct=66697
676. Baker TB, Brandon TH, Chassin L. Motivational influences on cigarette smoking. Annu Rev Psychol 2004;55:463-91.
677. Jackson C, Brown JD, L'Engle KL. R-rated movies, bedroom televisions, and initiation of smoking by white and black adolescents. Arch Pediatr Adolesc Med 2007;161:260-8.
678. Lovato C, Linn G, Stead LF, et al. Impact of tobacco advertising and promotion on increasing adolescent smoking behaviours. Cochrane Database Syst Rev 2003:CD003439.
679. McCool JP, Cameron LD, Petrie KJ. The

700. Health Research Inc: National Youth

assessment of abuse liability. Psychopharmacology (Berl) 2003;167:20-7.
729. Moolchan ET, Robinson ML, Ernst M, et al. Safety and efficacy of the nicotine patch and gum for the treatment of adolescent tobacco addiction. Pediatrics 2005;115:e407-14.
730. Niederhofer H, Huber M. Bupropion may support psychosocial treatment of nicotine-dependent adolescents: preliminary results. Pharmacotherapy 2004;24:1524-8.
731. Muramoto ML, Leischow SJ, Sherrill D, et al. Randomized, double-blind, plac-2(b3pe)5(ndent a)ch .LrmzJ g9(Nied)6(erh)6[(e)ntrollei3T, -0a5(o)6

2000. Available at: http://cancerweb.ncl.ac.uk/cancernet/600365.html.
756. Streppel MT, Boshuizen HC, Ocke MC, et al. Mortality and life expectancy in relation to long-term cigarette, cigar and pipe smoking: the Zutphen Study. Tob Control 2007;16:107-13.
757. Maxwell JC. Cigar industry in 2005. Richmond, VA, The Maxwell Report, 2006,
758. Delnevo CD, Hrywna M. “A Whole ‘Nother Smoke” or a Cigarette in Disguise: How RJ Reynolds Reframed the Image of Little Cigars. Am J Public Health 2007; 97:1368-75.
759. Nyman AL, Taylor TM, Biener L. Trends in cigar smoking and perceptions of health risks among Massachusetts adults. Tob Control 2002;11 Suppl 2:ii25-8.
760. International Agency for Research on Cancer. Tobacco smoke and involuntary smoking. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans. Lyon, International Agency for Research on Cancer, 2004.
761. Wetter DW, McClure JB, de Moor C, et al. Concomitant use of cigarettes and smokeless tobacco: prevalence, correlates, and predictors of tobacco cessation. Prev Med 2002;34:638-48.
762. Tomar SL. Is use of smokeless tobacco a risk factor for cigarette smoking? The U.S.

781. England LJ, Grauman A, Qian C, et al. Misclassification of maternal smoking status and its effects on an epidemiologic study of pregnancy outcomes. Nicotine Tob Res 2007;9:1005-13.
782. Webb DA, Boyd NR, Messina D, et al. The discrepancy between self-reported smoking status and urine continine levels among women enrolled in prenatal care at four publicly funded clinical sites. J Public Health Manag Pract 2003;9:322-5.
783. Mullen PD, Carbonari JP, Tabak ER, et al. Improving disclosure of smoking by pregnant women. Am J Obstet Gynecol 1991;165:409-13.
784. Lindqvist R, Lendahls L, Tollbom O, et al. Smoking during pregnancy: comparison of self-reports and cotinine levels in 496 women. Acta Obstet Gynecol Scand 2002;81:240-4.
785. England LJ, Kendrick JS, Wilson HG, et al. Effects of smoking reduction during pregnancy on the birth weight of term infants. Am J Epidemiol 2001;154:694-701.
786. Lieberman E, Gremy I, Lang JM, et al. Low birthweight at term and the timing of fetal exposure to maternal smoking. Am J Public Health 1994;84:1127-31.
787. DiClemente CC, Dolan-Mullen P, Windsor RA. The process of pregnancy smoking cessation: implications for interventions. Tob Control 2000;9 Suppl 3:III16-21.
788. U.S. Department of Health and Human Services. Women and smoking: A Report of the Surgeon General. Atlanta, GA, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2001.
789. American Society for Reproductive Medicine. Smoking and infertility. Available at: http://www.asrm.org/Patients/FactSheets/smoking.pdf.
790. Smoking and infertility. Fertil Steril 2006;86:S172-7.
791. Colman GJ, Joyce T. Trends in smoking before, during, and after pregnancy in ten states. Am J Prev Med 2003;24:29-35.
792. Lelong N, Kaminski M, Saurel-Cubizolles MJ, et al. Postpartum return to smoking among usual smokers who quit during pregnancy. Eur J Public Health 2001;11:334-9.
793. Matturri L, Ottaviani G, Lavezzi AM. Maternal smoking and sudden infant death syndrome: epidemiological study related to pathology. Virchows Arch 2006;449:697-706.
794. Lannero E, Wickman M, Pershagen G, et al. Maternal smoking during pregnancy increases the risk of recurrent wheezing during the first years of life (BAMSE). Respir Res 2006;7:3.
795. Billaud N. [What are the other long-term consequences of maternal smoking during pregnancy?]. J Gynecol Obstet Biol Reprod (Paris) 2005;34 Spec No 1:3S234-40.
796. Shenassa ED, Brown MJ. Maternal smoking and infantile gastrointestinal dysregulation: the case of colic. Pediatrics 2004;114:e497-505.
797. Schmitz M, Denardin D, Laufer Silva T, et al. Smoking during pregnancy and attention-deficit/hyperactivity disorder, predominantly inattentive type: a case-control study. J Am Acad Child Adolesc Psychiatry 2006;45:1338-45.
798. Vielwerth SE, Jensen RB, Larsen T, et al. The impact of maternal smoking on fetal and infant growth. Early Hum Dev 2007;83:491-5.
799. French GM, Groner JA, Wewers ME, et al. Staying smoke free: an intervention to prevent postpartum relapse. Nicotine Tob Res 2007;9:663-70.
800. Windsor RA, Woodby LL, Miller TM, et al. Effectiveness of Agency for Health Care Policy and Research clinical practice guideline and patient education methods for pregnant smokers in Medicaid maternity care. Am J Obstet Gynecol 2000;182:68-75.
801. Wisborg K, Henriksen TB, Jespersen LB, et al. Nicotine patches for pregnant smokers: a randomized controlled study. Obstet Gynecol 2000;96:967-71.
802. Kapur B, Hackman R, Selby P, et al. Randomized, double-blind, placebo-controlled trial of nicotine replacement therapy in pregnancy. Curr Ther Res Clin Exp 2001;62:274-8.
803. Pollack KI, Oncken C, Lipkus IM, et al. Nicotine replacement and behavioral therapy for smoking cessation in pregnancy. Am J Prev Med. 2007 Oct;33(4):297-305.
804. Armour BS, Campbell VA, Crews JE, et al. State-level prevalence of cigarette smoking and treatment advice, by disability status, United States, 2004. Prev Chronic Dis 2007;4:A86.
805. Nielsen A, Hannibal CG, Lindekilde BE. Maternal smoking predicts the risk of spontaneous abortion. Acta Obstet Gynecol Scand 2006;85:1057-65.
806. Kyrklund-Blomberg NB, Granath F, Cnattingius S. Maternal smoking and causes

831. Vander Weg MW, Klesges RC, Ward KD. Differences in resting energy expenditure between black and white smokers: implications for postcessation weight gain. Eur J Clin Nutr 2000;54:895-9.
832. Borrelli B, Spring B, Niaura R, et al. Influences of gender and weight gain on short-term relapse to smoking in a cessation trial. J Consult Clin Psychol 2001;69:511-5.
833. Delnevo CD, Hrywna M, Abatemarco DJ, et al. Relationships between cigarette smoking and weight control in young women. Fam Community Health 2003;26:140-6.
834. Meyers AW, Klesges RC, Winders SE, et al. Are weight concerns predictive of smoking cessation? A prospective analysis. J Consult Clin Psychol 1997;65:448-52.
835. Plotnikoff RC, Bercovitz K, Rhodes RE, et al. Testing a conceptual model related to weight perceptions, physical activity and smoking in adolescents. Health Educ Res 2007;22:192-202.
836. Pomerleau CS, Zucker AN, Stewart AJ. Characterizing concerns about post-cessation weight gain: results from a national survey of women smokers. Nicotine Tob Res 2001;3:51-60.
837. Vidrine JI, Anderson CB, Pollak KI, et al. Gender differences in adolescent smoking: mediator and moderator effects of self-generated expected smoking outcomes. Am? A3rcrm]TJns. J Co:cb3ferenPres Tt39:53(71(:448)5(-52.)]TJEMC 7/P <<

856. Centers for Disease Control and Prevention. Physical activity and health. 2005. Available at: http://www.cdc.gov/nccdphp/sgr/chap1.htm.